# Interpretation of Angular Distributions of Z-boson Production

Jen-Chieh Peng

University of Illinois at Urbana-Champaign

"QCD Evolution 2016" Workshop Nikhef, Amsterdam May 30 – June 3, 2016

Based on the paper of JCP, Wen-Chen Chang, Evan McClellan, Oleg Teryaev, Phys. Lett. B758 (2016) 384, arXiv: 1511.08932

#### **The Drell-Yan Process**

#### MASSIVE LEPTON-PAIR PRODUCTION IN HADRON-HADRON COLLISIONS AT HIGH ENERGIES\*

Sidney D. Drell and Tung-Mow Yan

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305 (Received 25 May 1970)

On the basis of a parton model studied earlier we consider the production process of large-mass lepton pairs from hadron-hadron inelastic collisions in the limiting region,  $s \rightarrow \infty$ ,  $Q^2/s$  finite,  $Q^2$  and s being the squared invariant masses of the lepton pair and the two initial hadrons, respectively. General scaling properties and connections with deep inelastic electron scattering are discussed. In particular, a rapidly decreasing cross section as  $Q^2/s \rightarrow 1$  is predicted as a consequence of the observed rapid falloff of the inelastic scattering structure function  $\nu W_2$  near threshold.



# Success and difficulties of the "naïve" Drell-YanSuccess:(T.M. Yan, hep-ph/9810268)

- Scaling of the cross sections (depends on x1 and x2 only)
- Nuclear dependence (cross section depends linearly on the mass A)
- Angular distributions (1+cos<sup>2</sup>  $\Theta$  distributions)

**Difficulties:** 

- Absolute cross sections (K-factor is needed)
- Transverse momentum distributions (much larger <p<sub>T</sub>> than expected)

# **Drell-Yan angular distribution** Lepton Angular Distribution of "naïve" Drell-Yan:

$$\frac{d\sigma}{d\Omega} = \sigma_0 (1 + \lambda \cos^2 \theta); \quad \lambda = 1$$



#### Data from Fermilab E772

(Ann. Rev. Nucl. Part. Sci. 49 (1999) 217-253)

## **Drell-Yan lepton angular distributions**



Θ and Φ are the decay polar and azimuthal angles of the  $μ^$ in the dilepton rest-frame

#### **Collins-Soper frame**

A general expression for Drell-Yan decay angular distributions:  $\left(\frac{1}{\sigma}\right)\left(\frac{d\sigma}{d\Omega}\right) = \left[\frac{3}{4\pi}\right]\left[1 + \lambda\cos^2\theta + \mu\sin2\theta\cos\phi + \frac{\nu}{2}\sin^2\theta\cos2\phi\right]$ Lam-Tung relation:  $1 - \lambda = 2\nu$ 

- Reflect the spin-1/2 nature of quarks
  (analog of the Callan-Gross relation in DIS)
- Insensitive to QCD corrections



 $v \neq 0$  and v increases with  $p_T$ 



Data from NA10 (Z. Phys. 37 (1988) 545)

Violation of the Lam-Tung relation suggests interesting new origins (Brandenburg, Nachtmann, Mirkes, Brodsky, Khoze, Müller, Eskolar, Hoyer, Väntinnen, Vogt, etc.)

#### Boer-Mulders function $h_1^{\perp}$ $\bigcirc$ - $\bigcirc$

- Boer pointed out that the cos2¢ dependence can be caused by the presence of the Boer-Mulders function.
- $h_1^{\perp}$  can lead to an azimuthal dependence with  $v \propto \left(\frac{h_1^{\perp}}{f_1}\right) \left(\frac{h_1^{\perp}}{\overline{f_1}}\right)$



Boer, PRD 60 (1999) 014012

 $h_{1}^{\perp}(x,k_{T}^{2}) = \frac{\alpha_{T}}{\pi} c_{H} \frac{M_{C}M_{H}}{k_{T}^{2} + M_{C}^{2}} e^{-\alpha_{T}k_{T}^{2}} f_{1}(x)$ 

$$v = 16\kappa_1 \frac{Q_T^2 M_C^2}{(Q_T^2 + 4M_C^2)^2}$$

$$\kappa_1 = 0.47, M_C = 2.3 \text{ GeV}$$

v>0 implies valence BM functions for pion and nucleon have same signs <sup>8</sup>



With Boer-Mulders function  $h_1^{\perp}$ :

 $v(\pi W \rightarrow \mu^{+} \mu^{-} X) \sim [valence h_{1}^{\perp}(\pi)] * [valence h_{1}^{\perp}(p)]$ 

 $v(pd \rightarrow \mu + \mu - X) \sim [valence h_1^{\perp}(p)] * [sea h_1^{\perp}(p)]$ 

Sea-quark BM function is much smaller than valence BM function



- Strong  $p_{T}\left(q_{T}\right)$  dependence of  $\lambda$  and  $\nu$
- Lam-Tung relation  $(1-\lambda = 2\nu)$  is satisfied within experimental uncertainties

## Recent CMS data for Z-boson production in p+p collision at 8 TeV



- Striking  $q_T$  dependencies for  $\lambda$  and  $\nu$  were observed at two rapidity regions
- Is Lam-Tung relation violated?

## Recent data from CMS for Z-boson production in p+p collision at 8 TeV



- Yes, the Lam-Tung relation is violated  $(1-\lambda > 2\nu)!$
- Can one understand the origin of the violation of the Lam-Tung relation?

Interpretation of the CMS Z-production results

$$\frac{d\sigma}{d\Omega} \propto (1 + \cos^2 \theta) + \frac{A_0}{2} (1 - 3\cos^2 \theta) + A_1 \sin 2\theta \cos \phi$$
$$+ \frac{A_2}{2} \sin^2 \theta \cos 2\phi + A_3 \sin \theta \cos \phi + A_4 \cos \theta$$
$$+ A_5 \sin^2 \theta \sin 2\phi + A_6 \sin 2\theta \sin \phi + A_7 \sin \theta \sin \phi$$

#### **Questions:**

- How is the above expression derived?
- Can one express  $A_0 A_7$  in terms of some quantities?
- Can one understand the  $Q_T$  dependence of  $A_0, A_1, A_2$ , etc?
- Can one understand the origin of the violation of Lam-Tung relation?

#### Define three planes in the Collins-Soper frame

1) Hadron Plane

- Contains the beam  $\vec{P}_B$  and target  $\vec{P}_T$  momenta
- Angle  $\beta$  satisfies the relation  $\tan \beta = q_T / Q$



#### Define three planes in the Collins-Soper frame

#### 1) Hadron Plane

- Contains the beam  $\vec{P}_B$  and target  $\vec{P}_T$  momenta
- Angle  $\beta$  satisfies the relation  $\tan \beta = q_T / Q$

#### 2) Quark Plane

- q and  $\overline{q}$  have head on collision along the  $\hat{z}'$  axis
- $\hat{z}'$  axis has angles  $\theta_1$  and  $\phi_1$  in the C S frame



#### Define three planes in the Collins-Soper frame

1) Hadron Plane

Φ

 $\vec{p}_B$ 

 $l^+$ 

61 Hadron Plane

 $\vec{p}_T$ 

epton Plane

 $\hat{y}$ 

 $\hat{z}$ 

 $\theta$ 

 $\hat{x}$ 

 $\theta_0$ 

Quark Plane

- Contains the beam  $\vec{P}_B$  and target  $\vec{P}_T$  momenta
- Angle  $\beta$  satisfies the relation  $\tan \beta = q_T / Q$

#### 2) Quark Plane

- q and  $\overline{q}$  have head on collision along the  $\hat{z}'$  axis
- $\hat{z}'$  axis has angles  $\theta_1$  and  $\phi_1$  in the C S frame

#### 3) Lepton Plane

- $l^-$  and  $l^+$  are emitted back to back with equal  $|\vec{P}|$
- $l^-$  is emitted at angle  $\theta$  and  $\phi$  in the C S frame

Ø

 $\vec{p}_B$ 

 $l^+$ 

 $\vec{p}_T$ 

 $\phi_1$  Hadron Plane

Lepton Plane

 $\hat{y}$ 

 $\hat{z}$ 

 $\theta$ 

 $\hat{x}$ 

 $\theta_0$ 

<u>What is the lepton angular distribution</u> with respect to the  $\hat{z}'$  (natural) axis?

$$\frac{d\sigma}{d\Omega} \propto 1 + \frac{a\cos\theta_0}{\cos\theta_0} + \cos^2\theta_0$$

How to express the angular distribution in terms of  $\theta$  and  $\phi$ ?

 $\cos\theta_0 = \cos\theta\cos\theta_1 + \sin\theta\sin\theta_1\cos(\phi - \phi_1)$ 



$$\frac{d\sigma}{d\Omega} \propto (1 + \cos^2 \theta) + \frac{\sin^2 \theta_1}{2} (1 - 3\cos^2 \theta) + (\frac{1}{2}\sin 2\theta_1 \cos \phi_1) \sin 2\theta \cos \phi + (\frac{1}{2}\sin^2 \theta_1 \cos 2\phi_1) \sin^2 \theta \cos 2\phi + (a\sin \theta_1 \cos \phi_1) \sin \theta \cos \phi + (a\cos \theta_1) \cos \theta + (\frac{1}{2}\sin^2 \theta_1 \sin 2\phi_1) \sin^2 \theta \sin 2\phi + (\frac{1}{2}\sin 2\theta_1 \sin \phi_1) \sin 2\theta \sin \phi + (a\sin \theta_1 \sin \phi_1) \sin \theta \sin \phi.$$

$$\frac{d\sigma}{d\Omega} \propto (1 + \cos^2 \theta) + \frac{\sin^2 \theta_1}{2} (1 - 3\cos^2 \theta) + (\frac{1}{2}\sin 2\theta_1 \cos \phi_1) \sin 2\theta \cos \phi + (\frac{1}{2}\sin^2 \theta_1 \cos 2\phi_1) \sin^2 \theta \cos 2\phi + (a\sin \theta_1 \cos \phi_1) \sin \theta \cos \phi + (a\cos \theta_1) \cos \theta + (\frac{1}{2}\sin^2 \theta_1 \sin 2\phi_1) \sin^2 \theta \sin 2\phi + (\frac{1}{2}\sin 2\theta_1 \sin \phi_1) \sin 2\theta \sin \phi + (a\sin \theta_1 \sin \phi_1) \sin \theta \sin \phi.$$

$$\frac{d\sigma}{d\Omega} \propto (1 + \cos^2 \theta) + \frac{A_0}{2} (1 - 3\cos^2 \theta) + A_1 \sin 2\theta \cos \phi + \frac{A_2}{2} \sin^2 \theta \cos 2\phi + A_3 \sin \theta \cos \phi + A_4 \cos \theta + A_5 \sin^2 \theta \sin 2\phi + A_6 \sin 2\theta \sin \phi + A_7 \sin \theta \sin \phi$$

## $A_0 - A_7$ are entirely described by $\theta_1, \phi_1$ and a

Angular distribution coefficients  $A_0 - A_7$ 



 $A_0 = \langle \sin^2 \theta_1 \rangle$  $A_1 = \frac{1}{2} \left\langle \sin 2\theta_1 \cos \phi_1 \right\rangle$  $A_2 = \left\langle \sin^2 \theta_1 \cos 2\phi_1 \right\rangle$  $A_3 = a \langle \sin \theta_1 \cos \phi_1 \rangle$  $A_4 = a \left\langle \cos \theta_1 \right\rangle$  $A_5 = \frac{1}{2} \left\langle \sin^2 \theta_1 \sin 2\phi_1 \right\rangle$  $A_6 = \frac{1}{2} \left\langle \sin 2\theta_1 \sin \phi_1 \right\rangle$  $A_7 = a \langle \sin \theta_1 \sin \phi_1 \rangle$ 

## Some implications of the angular distribution coefficients $A_0 - A_7$

- $A_0 = \langle \sin^2 \theta_1 \rangle$  $A_1 = \frac{1}{2} \left\langle \sin 2\theta_1 \cos \phi_1 \right\rangle$  $A_2 = \left\langle \sin^2 \theta_1 \cos 2\phi_1 \right\rangle$  $A_3 = a \left\langle \sin \theta_1 \cos \phi_1 \right\rangle$  $A_{4} = a \langle \cos \theta_{1} \rangle$  $A_5 = \frac{1}{2} \left\langle \sin^2 \theta_1 \sin 2\phi_1 \right\rangle$  $A_6 = \frac{1}{2} \left\langle \sin 2\theta_1 \sin \phi_1 \right\rangle$  $A_7 = a \left\langle \sin \theta_1 \sin \phi_1 \right\rangle$
- $A_0 \ge A_2 \text{ (or } 1 \lambda 2\nu \ge 0)$
- Lam Tung relation  $(A_0 = A_2)$ is satisfied when  $\phi_1 = 0$
- Forward backward asymmetry,*a*, is reduced by a factor of  $\langle \cos \theta_1 \rangle$  for  $A_4$
- $A_5, A_6, A_7$  are odd function of  $\phi_1$  and must vanish from symmetry consideration
- Some equality and inequality relations among  $A_0 - A_7$  can be obtained 21





# Compare with CMS data on $\lambda$ (*Z* production in *p*+*p* collision at 8 TeV)



## Compare with CMS data on v (*Z* production in *p*+*p* collision at 8 TeV)



$$v = \frac{2q_T^2}{2Q^2 + 3q_T^2} \quad \text{for } q\overline{q} \to Zg$$
$$v = \frac{10q_T^2}{2Q^2 + 15q_T^2} \quad \text{for } qG \to Zq$$

Dashed curve corresponds to a mixture of 58.5% qGand 41.5%  $q\bar{q}$  processes

Solid curve corresponds to

$$\left\langle \sin^2 \theta_1 \cos 2\phi_1 \right\rangle / \left\langle \sin^2 \theta_1 \right\rangle = 0.77$$

 $q - \bar{q}$  axis is non-coplanar relative to the hadron plane <sub>25</sub>

Origins of the non-coplanarity 1) Processes at order  $\alpha_s^2$  or higher



2) Intrinsic  $k_T$  from inetracting partons

## Compare with CMS data on Lam-Tung relation



Solid curves correspond to a mixture of 58.5% qG and 41.5%  $q\overline{q}$  processes, and  $\langle \sin^2 \theta_1 \cos 2\phi_1 \rangle / \langle \sin^2 \theta_1 \rangle = 0.77$ 

## Violation of Lam-Tung relation is well described

#### Compare with CDF data (*Z* production in $p + \bar{p}$ collision at 1.96 TeV)



Solid curves correspond to a mixture of 27.5% qG and 72.5%  $q\overline{q}$  processes, and  $\langle \sin^2 \theta_1 \cos 2\phi_1 \rangle / \langle \sin^2 \theta_1 \rangle = 0.85$ 

Violation of Lam-Tung relation is not ruled out

# Summary

- The lepton angular distribution coefficients  $A_0$ - $A_7$  are described in terms of the polar and azimuthal angles of the  $q \bar{q}$  axis.
- The striking  $q_T$  dependence of  $A_0$  (or equivalently,  $\lambda$ ) can be well described by the mis-alignment of the  $q \bar{q}$  axis and the Collins-Soper *z*-axis.
- Violation of the Lam-Tung relation  $(A_0 \neq A_2)$  is described by the non-coplanarity of the  $q - \overline{q}$ axis and the hadron plane. This can come from order  $\alpha_s^2$  or higher processes or from intrinsic  $k_T$ .
- This study can be extended to fixed-target Drell-Yan data.