Gluon TMDs in the small-x limit



Tom van Daal

Collaborators: D. Boer, S. Cotogno, P.J. Mulders, A. Signori, Y. Zhou



QCD Evolution 2016, Amsterdam





- Introduction
- Gluon TMDs for various target polarizations
- Gluon TMDs in the small-x limit
- Conclusion

What is a TMD?

- A hadron correlator is parametrized in terms of transverse momentum dependent (TMD) parton distribution functions (PDFs), also called TMDs.
- A TMD is a density function in the longitudinal momentum fraction *x* and the transverse momentum *k*_τ, encoding the 3D internal structure of hadrons.



parton momentum:

 $k^{\mu} = xP^{\mu} + k^{\mu}_{\tau} + \sigma n^{\mu}$

Vector and tensor polarized targets

Density matrix for spin-1 targets:



Parametrizing TMD distribution correlators

extracted parton

		quark	gluon
ol.	unpolarized		
get p	vector polarized		
taı	tensor polarized		×

[Tangerman, Mulders 1994; Boer, Mulders 1997; Bacchetta, Mulders 2000; Mulders, Rodrigues 2001]

The gluon-gluon TMD correlator





- Parametrization in terms of Lorentz structures
- Respecting hermiticity and invariance under parity
- T-odd functions are allowed

The gluon-gluon TMD correlator



$$\Gamma^{ij}(x,\boldsymbol{k}_{T}) \equiv \int \left. \frac{d\xi \cdot P \, d^{2}\xi_{T}}{(2\pi)^{3}} \, e^{ik \cdot \xi} \left\langle P \right| F^{ni}(0) \boldsymbol{U}_{[0,\xi]} F^{nj}(\xi) \boldsymbol{U}_{[\xi,0]}' \left| P \right\rangle \right|_{\xi \cdot n = 0}$$



process-dependent gauge links

[Mulders, Rodrigues 2001]

gluon pol.

- Parametrization in terms of Lorentz structures
- Respecting hermiticity and invariance under parity
- T-odd functions are allowed

	Gluons	$-g_{\scriptscriptstyle T}^{ij}$	$i\epsilon_{\scriptscriptstyle T}^{ij}$	$k_T^i, k_T^{ij}, ext{etc.}$	
	U	f_1		h_1^\perp	
<u>ol</u> .	L		g_1	h_{1L}^{\perp}	
et p	Т	f_{1T}^{\perp}	g_{1T}	h_1,h_{1T}^\perp	
rge	LL	f_{1LL}		h_{1LL}^{\perp}	
ta	LT	f_{1LT}	g_{1LT}	h_{1LT}, h_{1LT}^{\perp}	
	TT	f_{1TT}	g_{1TT}	$\boldsymbol{h_{1TT}},h_{1TT}^{\perp},h_{1TT}^{\perp\perp}$	

Definite rank gluon TMDs

Lorentz expansion in terms of (leading twist) definite rank gluon TMDs:

$$\begin{split} \Gamma_{U}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-g_{T}^{ij} f_{1}(x, \mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij}}{M^{2}} h_{1}^{\perp}(x, \mathbf{k}_{T}^{2}) \right], \\ \Gamma_{L}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[i \epsilon_{T}^{ij} S_{L} g_{1}(x, \mathbf{k}_{T}^{2}) + \frac{\epsilon_{T}^{ii} \alpha k_{T}^{ij} \alpha S_{L}}{2M^{2}} h_{1L}^{\perp}(x, \mathbf{k}_{T}^{2}) \right], \\ \Gamma_{T}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \epsilon_{T}^{STk_{T}}}{M} f_{1T}^{\perp}(x, \mathbf{k}_{T}^{2}) + \frac{i \epsilon_{T}^{ij} k_{T} \cdot S_{T}}{M} g_{1T}(x, \mathbf{k}_{T}^{2}) \\ &- \frac{\epsilon_{T}^{k_{T}}(s_{T}^{j}) + \epsilon_{T}^{S}(k_{T}^{j})}{4M} h_{1}(x, \mathbf{k}_{T}^{2}) - \frac{\epsilon_{T}^{ii} \alpha k_{T}^{j} \alpha S_{T}}{2M^{3}} h_{1T}^{\perp}(x, \mathbf{k}_{T}^{2}) \\ &- \frac{\epsilon_{T}^{ij} S_{LL} f_{1LL}(x, \mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij} S_{LL}}{M^{2}} h_{1LL}(x, \mathbf{k}_{T}^{2}) \\ &- \frac{\epsilon_{T}^{ij} s_{LT} f_{1LT}(x, \mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij} S_{LL}}{M^{2}} h_{1LL}^{\perp}(x, \mathbf{k}_{T}^{2}) \right], \end{split}$$

$$\Gamma_{LT}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-g_{T}^{ij} \mathbf{k}_{T} \cdot S_{LT} f_{1LT}(x, \mathbf{k}_{T}^{2}) + \frac{i \epsilon_{T}^{ij} \epsilon_{T} S_{LT} k_{T}}{M^{2}} h_{1LT}^{\perp}(x, \mathbf{k}_{T}^{2}) \right], \\ \Gamma_{LT}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \mathbf{k}_{T} \cdot S_{LT}}{M} f_{1LT}(x, \mathbf{k}_{T}^{2}) + \frac{i \epsilon_{T}^{ij} \epsilon_{T} S_{LT} k_{T}}{M} g_{1LT}(x, \mathbf{k}_{T}^{2}) \right], \\ \Gamma_{TT}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \mathbf{k}_{T} \cdot S_{TT}}{M} f_{1LT}(x, \mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij} \alpha S_{LT} \alpha}{M^{3}} h_{1LT}^{\perp}(x, \mathbf{k}_{T}^{2}) \right], \\ \Gamma_{TT}^{ij}(x, \mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} k_{T} \alpha \beta S_{TT} \alpha \beta}{M^{2}} f_{1TT}(x, \mathbf{k}_{T}^{2}) + \frac{i \epsilon_{T}^{ij} \epsilon_{T} \alpha k_{T}^{ij} \alpha S_{TT} \alpha \beta}{M^{2}} g_{1TT}(x, \mathbf{k}_{T}^{2}) \right] \right] \end{cases}$$

Constant tensors:

- $g_T^{\mu\nu} \equiv g^{\mu\nu} P^{\{\mu} n^{\nu\}}$
- $\epsilon^{\mu\nu}_{\scriptscriptstyle T} \equiv \epsilon^{Pn\mu\nu}$

Symmetric traceless tensors:

$$\begin{split} k_{T}^{ij} &\equiv k_{T}^{i} k_{T}^{j} + \frac{1}{2} \mathbf{k}_{T}^{2} g_{T}^{ij}, \\ k_{T}^{ijk} &\equiv k_{T}^{i} k_{T}^{j} k_{T}^{k} + \frac{1}{4} \mathbf{k}_{T}^{2} \left(g_{T}^{ij} k_{T}^{k} + g_{T}^{ik} k_{T}^{j} + g_{T}^{jk} k_{T}^{i} \right) \\ & \dots \text{ only 2 independent components:} \end{split}$$

 $k_{\scriptscriptstyle T}^{i_1...i_n} \quad \longleftrightarrow \quad |m{k}_{\scriptscriptstyle T}|^n \, e^{inarphi}$

Advantage of definite rank TMDs:

 One-to-one mapping between TMDs in *k*_T and *b*_T space (relevant for evolution)

Definite rank gluon TMDs

Lorentz expansion in terms of (leading twist) definite rank gluon TMDs:

$$\begin{split} \Gamma_{U}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-g_{T}^{ij} f_{1}(x,\mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij}}{M^{2}} h_{T}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{L}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[i\epsilon_{T}^{ij} S_{L} g_{1}(x,\mathbf{k}_{T}^{2}) + \frac{\epsilon_{T}^{ij} (k_{T}^{ij})^{\alpha} S_{L}}{2M^{2}} h_{1L}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{T}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \epsilon_{T}^{STkT}}{M} f_{1T}^{\perp}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \mathbf{k}_{T} \cdot \mathbf{S}_{T}}{M} g_{1T}(x,\mathbf{k}_{T}^{2}) \\ &- \frac{\epsilon_{T}^{kT}(iS_{T}^{ij}) + \epsilon_{T}^{ST}(ik_{T}^{ij})}{4M} h_{1}(x,\mathbf{k}_{T}^{2}) - \frac{\epsilon_{T}^{ij} (k_{T}^{ij} k_{T} \cdot \mathbf{S}_{T}}{2M^{3}} h_{1T}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{LL}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-g_{T}^{ij} S_{LL} f_{1LL}(x,\mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij} S_{LL}}{M^{2}} h_{1LL}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{LT}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-g_{T}^{ij} K_{LT} f_{1LT}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T}^{SLTkT}}{M^{2}} h_{1LT}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{LT}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \mathbf{k}_{T} \cdot \mathbf{S}_{LT}}{M} f_{1LT}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T}^{SLTkT}}{M^{3}} h_{1LT}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{LT}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} \mathbf{k}_{T} \cdot \mathbf{S}_{LT}}{M} f_{1LT}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T}^{SLTkT}}{M^{3}} h_{1LT}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{TT}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} k_{T} \cdot \mathbf{S}_{TT\alpha\beta}}{M^{2}} f_{1TT}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T}^{\beta} k_{T}^{\gamma} \cdot \mathbf{S}_{TT\alpha\beta}}{M^{3}} h_{1LT}^{\perp}(x,\mathbf{k}_{T}^{2}) \right], \\ \Gamma_{TT}^{ij}(x,\mathbf{k}_{T}) &= \frac{x}{2} \left[-\frac{g_{T}^{ij} k_{T} \cdot \mathbf{S}_{T\alpha\beta}}{M^{2}} f_{1TT}(x,\mathbf{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T}^{\beta} k_{T}^{\gamma} \cdot \mathbf{S}_{TT\alpha\beta}}{M^{2}} h_{1TT}^{1}(x,\mathbf{k}_{T}^{2}) + \frac{k_{T}^{ij} \epsilon_{T}^{\beta} k_{T}^{\gamma} \cdot \mathbf{S}_{T\alpha\beta}}{M^{2}} h_{1TT}^{\beta}(x,\mathbf{k}_{T}^{2}) \right], \\ \end{array}$$

Constant tensors:

- $g_T^{\mu\nu} \equiv g^{\mu\nu} P^{\{\mu} n^{\nu\}}$
- $\epsilon^{\mu\nu}_{\scriptscriptstyle T} \equiv \epsilon^{Pn\mu\nu}$

Symmetric traceless tensors:

$$\begin{aligned} k_{T}^{ij} &\equiv k_{T}^{i} k_{T}^{j} + \frac{1}{2} \boldsymbol{k}_{T}^{2} g_{T}^{ij}, \\ k_{T}^{ijk} &\equiv k_{T}^{i} k_{T}^{j} k_{T}^{k} + \frac{1}{4} \boldsymbol{k}_{T}^{2} \left(g_{T}^{ij} k_{T}^{k} + g_{T}^{ik} k_{T}^{j} + g_{T}^{jk} k_{T}^{i} \right) \end{aligned}$$

... only 2 independent components: $k_{\tau}^{i_1...i_n} \iff |\mathbf{k}_{\tau}|^n e^{in\varphi}$

Advantage of definite rank TMDs:

 One-to-one mapping between TMDs in *k*_T and *b*_T space (relevant for evolution)

Gluon vs. quark PDFs

[NNPDF Collaboration, 2012]



- Gluons dominate over quarks at small x
- What happens to the gluon TMDs as $x \rightarrow 0$?

The gluon-gluon correlator at x=0

$$\begin{split} \Gamma^{[+,-]\,ij}(0,\boldsymbol{k}_{T}) &\equiv \int \frac{d\xi \cdot P \, d^{2}\xi_{T}}{(2\pi)^{3}} \, e^{ik \cdot \xi} \left\langle P \right| F^{ni}(0) \, U_{[0,\xi]}^{[+]} \, F^{nj}(\xi) \, U_{[\xi,0]}^{[-]} \left| P \right\rangle \Big|_{\xi \cdot n = k \cdot n = 0} & \left[a \text{ "dipole-type" operator} \right] \\ &= \frac{k_{T}^{i} k_{T}^{j}}{2\pi L} \int \frac{d^{2}\xi_{T}}{(2\pi)^{2}} \, e^{ik_{T} \cdot \xi_{T}} \left\langle P \right| \left(U^{[\Box]} - I \right) \left| P \right\rangle \Big|_{\xi \cdot n = 0} \\ &\equiv \frac{k_{T}^{i} k_{T}^{j}}{2\pi L} \, \Gamma_{0}^{[\Box]}(\boldsymbol{k}_{T}) & \text{[the Wilson loop correlator, satisfying } \int d^{2}k_{T} \, \Gamma_{0}^{[\Box]}(\boldsymbol{k}_{T}) = 0 & \text{]} \end{split}$$



How to match f,g,h-type with e-type TMDs?

$\Gamma^{[+,-]ij}(x,\boldsymbol{k}_{T}) \equiv \int \frac{d\xi \cdot P d^{2}\xi_{T}}{(2\pi)^{3}} e^{ik\cdot\xi} \langle P F^{ni}(0) U^{[+]}_{[0,\xi]} F^{nj}(\xi) U^{[-]}_{[\xi,0]} P\rangle \bigg _{\xi\cdot n=0}$	$\Gamma_0^{[\Box]}(\boldsymbol{k}_T) \equiv \int \frac{d^2 \xi_T}{(2\pi)^2} e^{i k_T \cdot \xi_T} \left\langle P \right \left(U^{[\Box]} - I \right) \left P \right\rangle \Big _{\boldsymbol{\xi} \cdot \boldsymbol{n} = 0}$
$\Gamma_U^{[+,-]ij}(x,m{k}_{\scriptscriptstyle T}) = rac{x}{2} \left[-g_{\scriptscriptstyle T}^{ij}f_1(x,m{k}_{\scriptscriptstyle T}^2) + rac{k_{\scriptscriptstyle T}^{ij}}{M^2}h_1^\perp(x,m{k}_{\scriptscriptstyle T}^2) ight],$	$\Gamma^{[\Box]}_{0U}\left(oldsymbol{k}_{\scriptscriptstyle T} ight) = rac{\pi L}{M^2}e(oldsymbol{k}_{\scriptscriptstyle T}^2),$
$\Gamma_L^{[+,-]ij}(x,\boldsymbol{k}_{\scriptscriptstyle T}) = \frac{x}{2} \left[i \epsilon_{\scriptscriptstyle T}^{ij} S_L g_1(x,\boldsymbol{k}_{\scriptscriptstyle T}^2) + \frac{\epsilon_{\scriptscriptstyle T\alpha}^{\{i} \boldsymbol{k}_{\scriptscriptstyle T}^{j\}\alpha} S_L}{2M^2} h_{1L}^{\perp}(x,\boldsymbol{k}_{\scriptscriptstyle T}^2) \right],$	$\Gamma^{[\Box]}_{0L}(oldsymbol{k}_T)=0,$
$\begin{split} \Gamma_T^{[+,-]ij}(x,\boldsymbol{k}_T) &= \frac{x}{2} \left[-\frac{g_T^{ij} \epsilon_T^{S_T k_T}}{M} f_{1T}^{\perp}(x,\boldsymbol{k}_T^2) + \frac{i \epsilon_T^{ij} \boldsymbol{k}_T \cdot \boldsymbol{S}_T}{M} g_{1T}(x,\boldsymbol{k}_T^2) \right. \\ &\left \frac{\epsilon_T^{k_T \{i} S_T^{j\}} + \epsilon_T^{S_T \{i} k_T^{j\}}}{4M} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_{T\alpha}^{\{i} k_T^{j\}\alpha S_T}}{2M^3} h_{1T}^{\perp}(x,\boldsymbol{k}_T^2) \right], \end{split}$	$\Gamma_{0T}^{[\Box]}(\boldsymbol{k}_{T}) = \frac{\pi L}{M^{2}} \frac{\epsilon_{T}^{S_{T}k_{T}}}{M} e_{T}(\boldsymbol{k}_{T}^{2}),$
$\Gamma_{LL}^{[+,-]ij}(x,\boldsymbol{k}_{T}) = \frac{x}{2} \left[-g_{T}^{ij}S_{LL} f_{1LL}(x,\boldsymbol{k}_{T}^{2}) + \frac{k_{T}^{ij}S_{LL}}{M^{2}} h_{1LL}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right],$	$\Gamma^{[\Box]}_{0LL}\left(oldsymbol{k}_{T} ight) = rac{\pi L}{M^2} S_{LL} e_{LL}(oldsymbol{k}_{T}^2),$
$\Gamma_{LT}^{[+,-]ij}(x,\boldsymbol{k}_{T}) = \frac{x}{2} \left[-\frac{g_{T}^{ij}\boldsymbol{k}_{T} \cdot \boldsymbol{S}_{LT}}{M} f_{1LT}(x,\boldsymbol{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij}\epsilon_{T}^{S_{LT}\boldsymbol{k}_{T}}}{M} g_{1LT}(x,\boldsymbol{k}_{T}^{2}) \right]$ $S^{\{i\ L^{j}\}} = p^{ij\alpha}S_{TT}$	$\Gamma_{0LT}^{[\Box]}\left(\boldsymbol{k}_{T}\right) = \frac{\pi L}{M^{2}} \frac{\boldsymbol{k}_{T} \cdot \boldsymbol{S}_{LT}}{M} e_{LT}(\boldsymbol{k}_{T}^{2}),$
$+ \frac{S_{LT}\kappa_{T}}{M} h_{1LT}(x, \boldsymbol{k}_{T}^{2}) + \frac{\kappa_{T}}{M^{3}} \frac{S_{LT}}{h_{1LT}^{\alpha}} h_{1LT}^{\perp}(x, \boldsymbol{k}_{T}^{2}) \bigg],$ $\Gamma_{TT}^{[+,-]ij}(x, \boldsymbol{k}_{T}) = \frac{x}{2} \left[-\frac{g_{T}^{ij} k_{T}^{\alpha\beta} S_{TT\alpha\beta}}{M^{2}} f_{1TT}(x, \boldsymbol{k}_{T}^{2}) + \frac{i\epsilon_{T}^{ij} \epsilon_{T\gamma}^{\beta} k_{T}^{\gamma\alpha} S_{TT\alpha\beta}}{M^{2}} g_{1TT}(x, \boldsymbol{k}_{T}^{2}) \right]$	$\Gamma_{0TT}^{[\Box]}\left(\boldsymbol{k}_{T}\right) = \frac{\pi L}{M^{2}} \frac{k_{T}^{\alpha\beta} S_{TT\alpha\beta}}{M^{2}} e_{TT}(\boldsymbol{k}_{T}^{2})$
$+ S_{TT}^{ij} h_{1TT}(x, \boldsymbol{k}_{T}^{2}) + \frac{S_{TT\alpha}^{\{i} \boldsymbol{k}_{T}^{j\}\alpha}}{M^{2}} h_{1TT}^{\perp}(x, \boldsymbol{k}_{T}^{2}) + \frac{k_{T}^{ij\alpha\beta} S_{TT\alpha\beta}}{M^{4}} h_{1TT}^{\perp\perp}(x, \boldsymbol{k}_{T}^{2}) \right]$	

Unpolarized gluon TMDs at x=0

$$\Gamma^{[+,-]\,ij}(x,oldsymbol{k}_{\scriptscriptstyle T}) \quad \stackrel{x \to 0}{\longrightarrow} \quad rac{k_{\scriptscriptstyle T}^i k_{\scriptscriptstyle T}^j}{2\pi L} \,\Gamma_0^{[\Box]}(oldsymbol{k}_{\scriptscriptstyle T})$$

$$\frac{1}{x} \Gamma^{[+,-]\,ij}(x,\boldsymbol{k}_{T}) = \frac{1}{x} \frac{x}{2} \left[-g_{T}^{ij} f_{1}(x,\boldsymbol{k}_{T}^{2}) + \frac{k_{T}^{ij}}{M^{2}} h_{1}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right]$$

$$\xrightarrow{x \to 0} \quad \frac{k_T^i k_T^j}{2\pi L x} \Gamma_0^{[\Box]}(\boldsymbol{k}_T) = \frac{k_T^i k_T^j}{2\pi L x} \left[\frac{\pi L}{M^2} e(\boldsymbol{k}_T^2) \right]$$

$$= \frac{1}{2} \left[-g_T^{ij} \frac{k_T^2}{2M^2} \frac{e(k_T^2)}{x} + \frac{k_T^{ij}}{M^2} \frac{e(k_T^2)}{x} \right]$$

Unpolarized gluon TMDs at x=0

$$\Gamma^{[+,-]\,ij}(x,\boldsymbol{k}_{T}) \quad \stackrel{x\to 0}{\longrightarrow} \quad \frac{k_{T}^{i}k_{T}^{j}}{2\pi L}\,\Gamma_{0}^{[\Box]}(\boldsymbol{k}_{T})$$

$$\frac{1}{x} \Gamma^{[+,-]\,ij}(x,\boldsymbol{k}_{T}) = \frac{1}{2} \left[-g_{T}^{ij} f_{1}(x,\boldsymbol{k}_{T}^{2}) + \frac{k_{T}^{ij}}{M^{2}} h_{1}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right]$$

$$\stackrel{x \to 0}{\longrightarrow} \quad \frac{k_T^i k_T^j}{2\pi L x} \Gamma_0^{[\Box]}(\boldsymbol{k}_T) = \frac{k_T^i k_T^j}{2x M^2} e(\boldsymbol{k}_T^2)$$
$$= \quad \frac{1}{2} \left[-g_T^{ij} \frac{\boldsymbol{k}_T^2}{2M^2} \frac{e(\boldsymbol{k}_T^2)}{x} + \frac{k_T^{ij}}{M^2} \frac{e(\boldsymbol{k}_T^2)}{x} \right]$$

Conclusion:
$$f_1 = \frac{\boldsymbol{k}_T^2}{2M^2} h_1^\perp \rightarrow \frac{\boldsymbol{k}_T^2}{2xM^2} e(\boldsymbol{k}_T^2)$$

$$\begin{cases} \int dk_T^2 \, e(\boldsymbol{k}_T^2) = 0 \\ \int dk_T^2 \, \boldsymbol{k}_T^2 \, e(\boldsymbol{k}_T^2) \neq 0 \\ \Rightarrow \ e(\boldsymbol{k}_T^2) \text{ has a node in } \boldsymbol{k}_T^2 \end{cases}$$

Transversely polarized gluon TMDs at x=0

$$\Gamma^{[+,-]\,ij}(x,\boldsymbol{k}_{T}) \quad \stackrel{x\to 0}{\longrightarrow} \quad \frac{k_{T}^{i}k_{T}^{j}}{2\pi L}\,\Gamma_{0}^{[\Box]}(\boldsymbol{k}_{T})$$

$$\frac{1}{x} \Gamma_T^{[+,-]\,ij}(x,\boldsymbol{k}_T) = \frac{1}{x} \frac{x}{2} \left[-\frac{g_T^{ij} \epsilon_T^{S_T k_T}}{M} f_{1T}^{\perp}(x,\boldsymbol{k}_T^2) + \frac{i \epsilon_T^{ij} \boldsymbol{k}_T \cdot \boldsymbol{S}_T}{M} g_{1T}(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{k_T \{i} S_T^{j\}} + \epsilon_T^{S_T \{i} k_T^{j\}}}{4M} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{\{i} \alpha} k_T^{j\} \alpha S_T}{2M^3} h_{1T}^{\perp}(x,\boldsymbol{k}_T^2) \right]$$

$$\stackrel{x \to 0}{\longrightarrow} \quad \frac{k_T^i k_T^j}{2\pi L x} \Gamma_{0T}^{[\Box]}(\mathbf{k}_T) = \frac{k_T^i k_T^j}{2\pi L x} \left[\frac{\pi L}{M^2} \frac{\epsilon_T^{S_T k_T}}{M} e_T(\mathbf{k}_T^2) \right]$$

$$= \quad \frac{1}{2} \left[-\frac{g_T^{ij} \epsilon_T^{S_T k_T}}{M} \frac{\mathbf{k}_T^2}{2M^2} \frac{e_T(\mathbf{k}_T^2)}{x} - \frac{\epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} + \epsilon_T^{S_T \{i} k_T^{j\}}}{4M} \frac{\mathbf{k}_T^2}{2M^2} \frac{e_T(\mathbf{k}_T^2)}{x} - \frac{\epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} + \epsilon_T^{S_T \{i} k_T^{jj}\}}{2M^2} \frac{\mathbf{k}_T^2}{2M^2} \frac{e_T(\mathbf{k}_T^2)}{x} - \frac{\epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} + \epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij}}{4M} \frac{\mathbf{k}_T^2}{2M^2} \frac{e_T(\mathbf{k}_T^2)}{x} - \frac{\epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij} \epsilon_T^{ij}}{2M^3} \left(-\frac{e_T(\mathbf{k}_T^2)}{x} \right)$$

Transversely polarized gluon TMDs at x=0

$$\Gamma^{[+,-]\,ij}(x,\boldsymbol{k}_{T}) \quad \stackrel{x\to 0}{\longrightarrow} \quad \frac{k_{T}^{i}k_{T}^{j}}{2\pi L}\,\Gamma_{0}^{[\Box]}(\boldsymbol{k}_{T})$$

$$\frac{1}{x} \Gamma_T^{[+,-]\,ij}(x,\boldsymbol{k}_T) = \frac{1}{2} \left[-\frac{g_T^{ij} \epsilon_T^{S_T k_T}}{M} f_{1T}^{\perp}(x,\boldsymbol{k}_T^2) + \frac{i \epsilon_T^{ij} \boldsymbol{k}_T \cdot \boldsymbol{S}_T}{M} g_{1T}(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{k_T \{i\}} S_T^{j\}} + \epsilon_T^{k_T \{i\}} S_T^{j\}} + \epsilon_T^{k_T \{i\}} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{ij} \epsilon_T^{k_T \{i\}} S_T^{j\}} + \epsilon_T^{k_T \{i\}} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{ij} \epsilon_T^{k_T \{i\}} S_T^{j\}} + \epsilon_T^{k_T \{i\}} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{ij} \epsilon_T^{k_T \{i\}} S_T^{j\}} + \epsilon_T^{k_T \{i\}} h_1(x,\boldsymbol{k}_T^2) - \frac{\epsilon_T^{ij} \epsilon_T^{k_T \{i\}} h_1(x,\boldsymbol{k}_T^2)}{2M^3} h_{1T}^{\perp}(x,\boldsymbol{k}_T^2) \right]$$

$$\stackrel{x \to 0}{\longrightarrow} \quad \frac{k_T^i k_T^j}{2\pi L x} \Gamma_{0T}^{[\Box]}(\boldsymbol{k}_T) = \frac{k_T^i k_T^j}{2x M^2} \frac{\epsilon_T^{S_T k_T}}{M} e_T(\boldsymbol{k}_T^2)$$

$$= \quad \frac{1}{2} \left[-\frac{g_T^{ij} \epsilon_T^{S_T k_T}}{M} \frac{\boldsymbol{k}_T^2}{2M^2} \frac{e_T(\boldsymbol{k}_T^2)}{x} - \frac{\epsilon_T^{k_T \{i} S_T^{j\}} + \epsilon_T^{S_T \{i} k_T^{j\}}}{4M} \frac{\boldsymbol{k}_T^2}{2M^2} \frac{e_T(\boldsymbol{k}_T^2)}{x} - \frac{\epsilon_T^{\{i} \alpha k_T^{j\} \alpha S_T}}{2M^3} \left(-\frac{e_T(\boldsymbol{k}_T^2)}{x} \right) \right]$$

Conclusion:
$$f_{1T}^{\perp} = h_1 = -\frac{k_T^2}{2M^2} h_{1T}^{\perp} \to \frac{k_T^2}{2xM^2} e_T(k_T^2), \quad g_{1T} = 0$$

Shows up in $\Gamma_0^{[\Box]} - \Gamma_0^{[\Box^{\dagger}]}$, in agreement with the results in [Boer, G. Echevarría, Mulders, Zhou 2015]

Gluon TMDs at x=0

U	$f_1 = rac{oldsymbol{k}_T^2}{2M^2} h_1^\perp o rac{oldsymbol{k}_T^2}{2xM^2} e$
L	$g_1 = h_{1L}^\perp = 0$
Т	$f_{1T}^{\perp} = h_1 = -\frac{{oldsymbol{k}}_T^2}{2M^2} h_{1T}^{\perp} o rac{{oldsymbol{k}}_T^2}{2xM^2} e_T, g_{1T} = 0$
LL	$f_{1LL}=rac{oldsymbol{k}_T^2}{2M^2}h_{1LL}^\perp ightarrowrac{oldsymbol{k}_T^2}{2xM^2}e_{LL}$
LT	$f_{1LT} = h_{1LT} = -\frac{k_T^2}{4M^2} h_{1LT}^{\perp} \rightarrow \frac{k_T^2}{4xM^2} e_{LT}, g_{1LT} = 0$
TT	$f_{1TT} = \frac{2M^2}{3k_T^2} h_{1TT} = -\frac{1}{2} h_{1TT}^{\perp} = \frac{k_T^2}{6M^2} h_{1TT}^{\perp \perp} \to \frac{k_T^2}{6xM^2} e_{TT}, g_{1TT} = 0$

In the small-x limit we see that

- ... the gluon TMDs either vanish or become equal.
- ... the (non-vanishing) gluon TMDs are proportional to 1/x (up to resummed logs).



• New gluon TMDs for tensor polarized targets have been introduced.

	quark	gluon
unpolarized	 Image: A set of the set of the	 ✓
vector polarized	 Image: A set of the set of the	 ✓
tensor polarized	~	



• New gluon TMDs for tensor polarized targets have been introduced.

	quark	gluon
unpolarized	~	 ✓
vector polarized	1	1
tensor polarized		

- The gluon-gluon correlator at x=0 reduces to the F.T. of a Wilson loop.
- Also the Wilson loop correlator can be parametrized in terms of TMDs.
- In the small-x limit the picture of gluon TMDs becomes very simple: they either vanish or become equal.

Backup slides

The quark-quark TMD correlator

$$\Phi_{ij}(x, \mathbf{k}_T) \equiv \int \left. \frac{d\xi \cdot P \, d^2 \xi_T}{(2\pi)^3} \, e^{ik \cdot \xi} \left\langle P | \, \overline{\psi}_j(0) U_{[0,\xi]} \psi_i(\xi) \, | P \right\rangle \right|_{\xi \cdot n = 0}$$



process-dependent gauge link: $U_{[0,\xi]} \equiv \mathcal{P} \exp\left(-ig \int_0^{\xi} d\eta^{\mu} A_{\mu}(\eta)\right)$

- Parametrization in terms of Dirac structures
- 18 TMDs/ 4 collinear PDFs

quark pol.

$$\begin{split} \Phi_{\boldsymbol{U}}(x,\boldsymbol{k}_{T}) &= \left[f_{1}(x,\boldsymbol{k}_{T}^{2}) + \frac{i\boldsymbol{k}_{T}}{M} h_{1}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right] \frac{\boldsymbol{p}}{2} ,\\ \Phi_{\boldsymbol{L}}(x,\boldsymbol{k}_{T}) &= \left[\gamma^{5}S_{L} g_{1}(x,\boldsymbol{k}_{T}^{2}) + \frac{\gamma^{5}\boldsymbol{k}_{T}S_{L}}{M} h_{1L}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right] \frac{\boldsymbol{p}}{2} ,\\ \Phi_{\boldsymbol{T}}(x,\boldsymbol{k}_{T}) &= \left[\frac{\epsilon_{T}^{S_{T}\boldsymbol{k}_{T}}}{M} f_{1T}^{\perp}(x,\boldsymbol{k}_{T}^{2}) + \frac{\gamma^{5}\boldsymbol{k}_{T}\cdot\boldsymbol{S}_{T}}{M} g_{1T}(x,\boldsymbol{k}_{T}^{2}) \right. \\ &+ \gamma^{5}\boldsymbol{\beta}_{T} h_{1}(x,\boldsymbol{k}_{T}^{2}) - \frac{\gamma^{5}\gamma_{\nu}\boldsymbol{k}_{T}^{\nu\rho}S_{T\rho}}{M^{2}} h_{1T}^{\perp}(x,\boldsymbol{k}_{T}^{2}) \right] \frac{\boldsymbol{p}}{2} \end{split}$$

	Quarks	γ^+	$\gamma^+\gamma^5$	$i\sigma^{i+}\gamma^5$
	U	f_1		h_1^\perp
0.	L		g_1	h_{1L}^{\perp}
с С	Т	f_{1T}^{\perp}	g_{1T}	$oldsymbol{h_1}, h_{1T}^\perp$
nge	LL	f_{1LL}		h_{1LL}^{\perp}
ta	LT	f_{1LT}	g_{1LT}	h_{1LT},h_{1LT}^{\perp}
	TT	f_{1TT}	g_{1TT}	h_{1TT},h_{1TT}^{\perp}

[Tangerman, Mulders 1994; Boer, Mulders 1997; Bacchetta, Mulders 2000]

Comparing the quark and gluon TMDs

Quarks	γ^+	$\gamma^+\gamma^5$	$i\sigma^{i+}\gamma^5$
U	f_1		h_1^\perp
L		g_1	h_{1L}^{\perp}
Т	f_{1T}^{\perp}	g_{1T}	$oldsymbol{h_1},h_{1T}^\perp$
LL	f_{1LL}		h_{1LL}^{\perp}
LT	f_{1LT}	g_{1LT}	h_{1LT},h_{1LT}^{\perp}
TT	f_{1TT}	g_{1TT}	h_{1TT},h_{1TT}^{\perp}

Gluons	$-g_{\scriptscriptstyle T}^{ij}$	$i\epsilon_{\scriptscriptstyle T}^{ij}$	$k_T^i, k_T^{ij}, ext{etc.}$
U	f_1		h_1^\perp
L		g_1	h_{1L}^{\perp}
Т	f_{1T}^{\perp}	g_{1T}	h_1, h_{1T}^\perp
LL	f_{1LL}		h_{1LL}^{\perp}
LT	f_{1LT}	g_{1LT}	h_{1LT},h_{1LT}^{\perp}
TT	f_{1TT}	g_{1TT}	$\boldsymbol{h_{1TT}},h_{1TT}^{\perp},h_{1TT}^{\perp\perp}$