



Nikhef

8 May 2020

LHCb Highlights



Niels Tuning 

On the menu

- Introduction
 - Precision measurements
 - The LHCb physics menu
- Selection of dishes:
 - Recent highlights on CP violation
 - Recent highlights on Rare decays (*aka Flavour Anomalies*)

History of Flavour physics

GIM mechanism in $K^0 \rightarrow \mu\mu$

Weak Interactions with Lepton-Hadron Symmetry*


S. L. GLASHOW, J. ILIOPOULOS, AND L. MAIANI†
 Lyman Laboratory of Physics, Harvard University, Cambridge, Massachusetts 02139
 (Received 5 March 1970)

We propose a model of weak interactions in which the currents are constructed out of four basic quark fields and interact with a charged massive vector boson. We show, to all orders in perturbation theory, that the leading divergences do not violate any strong-interaction symmetry and the next to the leading divergences respect all observed weak-interaction selection rules. The model features a remarkable symmetry between leptons and quarks. The extension of our model to a complete Yang-Mills theory is discussed.

splitting, beginning at order $G(G\Lambda^2)$, as well as contributions to such unobserved decay modes as $K_2 \rightarrow \mu^+ + \mu^-$, $K^+ \rightarrow \pi^+ + l + \bar{l}$, etc., involving neutral lepton

We wish to propose a simple model in which the divergences are properly ordered. Our model is founded in a quark model, but one involving **four, not three,** fundamental fermions; the weak interactions are medi-

new quantum number C for charm.



Glashow, Iliopoulos, Maiani,
 Phys.Rev. D2 (1970) 1285

CP violation, $K_L^0 \rightarrow \pi\pi$

27 JULY 1964

EVIDENCE FOR THE 2π DECAY OF THE K_2^0 MESON*†

J. H. Christenson, J. W. Cronin,† V. L. Fitch,† and R. Turlay§
 Princeton University, Princeton, New Jersey
 (Received 10 July 1964)

This Letter reports the results of experimental studies designed to search for the 2π decay of the K_2^0 meson. Several previous experiments have

Progress of Theoretical Physics, Vol. 49, No. 2, February 1973

CP-Violation in the Renormalizable Theory of Weak Interaction

Makoto KOBAYASHI and Toshihide MASKAWA

Department of Physics, Kyoto University, Kyoto

(Received September 1, 1972)

doublet with the same charge assignment. This is because all phases of elements of a 3×3 unitary matrix cannot be absorbed into the phase convention of six fields. This possibility of CP -violation will be discussed later on.

Christenson, Cronin, Fitch, Turlay,
 Phys.Rev.Lett. 13 (1964) 138
 Kobayashi, Maskawa,
 Prog.Theor. Phys. 49 (1973) 652

$B^0 \leftrightarrow \bar{B}^0$ mixing

DESY 87-029
 April 1987

OBSERVATION OF $B^0 \cdot \bar{B}^0$ MIXING

The ARGUS Collaboration

In summary, the combined evidence of the investigation of B^0 meson pairs, lepton pairs and B^0 meson-lepton events on the $Y(4S)$ leads to the conclusion that $B^0 \cdot \bar{B}^0$ mixing has been observed and is substantial.

Parameters	Comments
$r > 0.09$ 90%CL	This experiment
$x > 0.44$	This experiment
$B^0 \cdot \bar{B}^0 \approx \tau_B < 160 \text{ MeV}$	B meson (\approx pion) decay constant
$m_b < 5 \text{ GeV}/c^2$	b-quark mass
$\tau_b < 1.4 \cdot 10^{-12} \text{ s}$	B meson lifetime
$ V_{td} < 0.018$	Kobayashi-Maskawa matrix element
$\eta_{CP} < 0.86$	QCD correction factor [17]
$m_t > 50 \text{ GeV}/c^2$	t quark mass

ARGUS Coll.
 Phys.Lett.B192 (1987) 245

Flavour physics has a track record

GIM mechanism in $K^0 \rightarrow \mu\mu$

Weak Interactions with Lepton-Hadron Symmetry*


S. L. GLASHOW, J. ILIOPoulos, AND L. MAIANI†
Lyman Laboratory of Physics, Harvard University, Cambridge, Massachusetts 02139
 (Received 5 March 1970)

We propose a model of weak interactions in which the currents are constructed out of four basic quark fields and interact with a charged massive vector boson. We show, to all orders in perturbation theory, that the leading divergences do not violate any strong-interaction symmetry and the next to the leading divergences respect all observed weak-interaction selection rules. The model features a remarkable symmetry between leptons and quarks. The extension of our model to a complete Yang-Mills theory is discussed.

splitting, beginning at order $G(G\Lambda^2)$, as well as contributions to such unobserved decay modes as $K_2 \rightarrow \mu^+ + \mu^-$, $K^+ \rightarrow \pi^+ + l + \bar{l}$, etc., involving neutral lepton

We wish to propose a simple model in which the divergences are properly ordered. Our model is founded in a quark model, but one involving four, not three, fundamental fermions; the weak interactions are medi-

new quantum number C for charm.



“...a quark model, but involving **four**, not three fundamental fermions...”

**Rare decay implied
 2nd up quark
 “discovery” of charm?**

CP violation, $K_L^0 \rightarrow \pi\pi$

27 JULY 1964

EVIDENCE FOR THE 2π DECAY OF THE K_2^0 MESON*†

J. H. Christenson, J. W. Cronin,† V. L. Fitch,† and R. Turlay§
Princeton University, Princeton, New Jersey
 (Received 10 July 1964)

This Letter reports the results of experimental studies designed to search for the 2π decay of the K_2^0 meson. Several previous experiments have

Progress of Theoretical Physics, Vol. 49, No. 2, February 1973

CP-Violation in the Renormalizable Theory of Weak Interaction

Makoto KOBAYASHI and Toshihide MASKAWA

Department of Physics, Kyoto University, Kyoto

(Received September 1, 1972)

doublet with the same charge assignment. This is because all phases of elements of a 3×3 unitary matrix cannot be absorbed into the phase convention of six fields. This possibility of CP-violation will be discussed later on.

“... phases of elements of 3×3 unitary matrix cannot be absorbed into [...] **six** fields ...”

**CP violation implied
 3rd family:
 “discovery” of bottom?**

$B^0 \leftrightarrow \bar{B}^0$ mixing

DESY 87-029
 April 1987

OBSERVATION OF $B^0 \cdot \bar{B}^0$ MIXING

The ARGUS Collaboration

In summary, the combined evidence of the investigation of B^0 meson pairs, lepton pairs and B^0 meson-lepton events on the $\Upsilon(4S)$ leads to the conclusion that $B^0 \cdot \bar{B}^0$ mixing has been observed and is substantial.

Parameters	Comments
$r > 0.09$ 90%CL	This experiment
$x > 0.44$	This experiment
$B^0 \cdot \bar{B}^0 \approx \tau_B < 160 \text{ MeV}$	B meson (\approx pion) decay constant
$m_b < 5 \text{ GeV}/c^2$	b-quark mass
$\tau_B < 1.4 \cdot 10^{-12} \text{ s}$	B meson lifetime
$ V_{td} < 0.018$	Kobayashi-Maskawa matrix element
$\eta_{\text{QCD}} < 0.86$	QCD correction factor [17]
$m_t > 50 \text{ GeV}/c^2$	t quark mass


“ $m_t > 50 \text{ GeV}/c^2$ t quark mass ”

**Mixing implied
 heavy quark:
 “discovery” of top?**

Precise flavour measurements








■ Historical record of indirect discoveries:


Particle	Indirect			Direct		
ν	β decay	Fermi	1932	Reactor ν -CC	Cowan, Reines	1956
W	β decay	Fermi	1932	$W \rightarrow e\nu$	UA1, UA2	1983
c	$K^0 \rightarrow \mu\mu$	GIM	1970	J/ψ	Richter, Ting	1974
b	CPV $K^0 \rightarrow \pi\pi$	CKM, 3 rd gen	1964/72	Y	Ledermann	1977
Z	ν -NC	Gargamelle	1973	$Z \rightarrow e^+e^-$	UA1	1983
t	B mixing	ARGUS	1987	$t \rightarrow Wb$	D0, CDF	1995
H	e^+e^-	EW fit, LEP	2000	$H \rightarrow 4\mu/\gamma\gamma$	CMS, ATLAS	2012
?	What's next ?		?			?



Precise flavour measurements


- Direct discoveries rightfully higher valued:

Particle	Indirect			Direct		
ν	β decay	Fermi	1932 	Reactor ν -CC	Cowan, Reines	1956 
W	β decay	Fermi	1932	$W \rightarrow e\nu$	UA1, UA2	1983 
c	$K^0 \rightarrow \mu\mu$	GIM	1970	J/ψ	Richter, Ting	1974 
b	CPV $K^0 \rightarrow \pi\pi$	CKM, 3 rd gen	1964/ 	Υ	Ledermann	1977
Z	ν -NC	Gargamelle	1973	$Z \rightarrow e^+e^-$	UA1	1983 
t	B mixing	ARGUS	1987	$t \rightarrow Wb$	D0, CDF	1995
H	e^+e^-	EW fit, LEP	2000	$H \rightarrow 4\mu/\gamma\gamma$	CMS, ATLAS	2012 
?	What's next ?		?			?



Precise flavour measurements

- Depending on your model, sensitive to multi-TeV scales, eg:



$$\mu_{B_s \rightarrow \mu^+ \mu^-} \simeq 1 \pm \frac{4\pi}{g^2 |V_{tb}^* V_{ts}|^2} \frac{v^2}{\Lambda^2}$$


$\mu_{B \rightarrow \mu\mu}$ is ratio $\text{BR}^{\text{exp}}/\text{BR}^{\text{SM}}$

$$\Lambda \gtrsim \frac{v}{\sqrt{0.2}} \times \begin{cases} \frac{\sqrt{4\pi}}{g |V_{tb}^* V_{ts}|} \\ 1 \end{cases} \simeq \begin{cases} 50 \text{ TeV}, & \text{anarchic tree} \\ 0.6 \text{ TeV}, & \text{MFV loop} \end{cases}$$

From Uli Haisch, [31 Aug 2016](#)
[arXiv:1510.03341](#)

Precise flavour measurements

- Depending on your model, sensitive to multi-TeV scales, eg:




From Uli Haisch, [31 Aug 2016](#)

On the menu


- Introduction
 - Precision measurements
 - The LHCb physics menu
- Selection of dishes:
 - Recent highlights on CP violation
 - Recent highlights on Rare decays (*aka Flavour Anomalies*)

LHCb Physics Landscape




LHCb Physics Landscape: more than b


Impressive $\sin^2\theta_W$




Resolve b and c jets



Anti-proton flux for cosmic rays




Improve proton pdf's




A.Garcia, R.Gauld,
A.Heijboer, J.Rojo
arXiv:2004.04756

Discovery of pentaquarks



CP violation in charm




$$A_{CP}(f; t) \equiv \frac{\Gamma(D^0(t) \rightarrow f) - \Gamma(\bar{D}^0(t) \rightarrow f)}{\Gamma(D^0(t) \rightarrow f) + \Gamma(\bar{D}^0(t) \rightarrow f)}$$

$$\Delta A_{CP} \equiv A_{CP}(K^- K^+) - A_{CP}(\pi^- \pi^+)$$

LHCb Physics Landscape: spectroscopy

Spectroscopy 35(29) hadrons discovered at LHC(b):



Collected by Patrick Koppenburg: <https://www.nikhef.nl/~pkoppenb/particles.html>

LHCb Physics Landscape

$b \rightarrow sll$

$B_s^0 \rightarrow \mu^+ \mu^-$
 R_K

$|V_{cb}|, |V_{ub}|$

$R(D)$

Today

CP violation:

γ

On the menu

- Introduction
 - Precision measurements
 - The LHCb physics menu
- Selection of dishes:
 - Recent highlights on CP violation
 - Recent highlights on Rare decays (*aka Flavour Anomalies*)

On the menu

- Introduction

- Precision measurements
- The LHCb physics menu

- Selection of dishes:

- Recent highlights on CP violation
- Recent highlights on Rare decays (*aka Flavour Anomalies*)


- New results

- 1) $|V_{cb}|$ with decay $B_s^0 \rightarrow D_s^* \bar{\mu}^+ \nu$
- 2) γ with decay $B^- \rightarrow D^0 (\rightarrow K_S^0 K^+ \pi^-) K^-$
- 3) γ with decay $B^0 \rightarrow D^0 K^{*0}$

- A remark on consistency


(CKM: a quick reminder...)

- 1) Matrix to transform weak- and mass-eigenstates:



(CKM: a quick reminder...)

- 1) Matrix to transform weak- and mass-eigenstates:




- 2) Matrix has imaginary numbers:

$$\begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}|e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}|e^{-i\beta} & -|V_{ts}|e^{i\beta_s} & |V_{tb}| \end{pmatrix}$$

(CKM: a quick reminder...)

- 1) Matrix to transform weak- and mass-eigenstates:




- 2) Matrix has imaginary numbers:

$$\begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}|e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}|e^{-i\beta} & -|V_{ts}|e^{i\beta_s} & |V_{tb}| \end{pmatrix}$$

- 3) Matrix is unitary:

$$V^+ V = \begin{pmatrix} V_{ud}^* & V_{cd}^* & V_{td}^* \\ V_{us}^* & V_{cs}^* & V_{ts}^* \\ V_{ub}^* & V_{cb}^* & V_{tb}^* \end{pmatrix} \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$V_{ub}^* V_{ud} + V_{cb}^* V_{cd} + V_{tb}^* V_{td} = 0$$



CKM: (1995) LHCb Letter-of-Intent

■ LHC-B Letter-of-Intent 1995





Figure 2.1: Limits on the CKM parameters (1σ) ρ and η for $m_t = 174 \text{ GeV}$. The annular region cen-

CKM: (1995) LHCb Letter-of-Intent ...

■ Letter-of-Intent 1995




Figure 2.1: Limits on the CKM parameters (1σ) ρ and η for $m_t = 174 \text{ GeV}$. The annular region cen-





Figure 2.2: The Unitarity Triangle




New measurement on $|V_{cb}|$

arXiv:2003.08453

- Measure decay rate of $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu$
 - Depends on momentum transfer q^2 :


$$\begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| e^{-i\beta} & -|V_{ts}| e^{i\beta_s} & |V_{tb}| \end{pmatrix}$$



$$\frac{d\Gamma(B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu)}{dq^2} = \frac{G_F^2 |V_{cb}|^2 |\eta_{EW}|^2 |\vec{p}| q^2}{96 \pi^3 m_{B^0}^2} \left(1 - \frac{m_\mu^2}{q^2}\right)^2$$

$$\times \left[(|H_+|^2 + |H_-|^2 + |H_0|^2) \left(1 + \frac{m_\mu^2}{2q^2}\right) + \frac{3}{2} \frac{m_\mu^2}{q^2} |H_t|^2 \right]$$

➤ Determine $|V_{cb}|$ and form factors



New measurement on $|V_{cb}|$

arXiv:2001.03225

- Measure rate relative to known B^0 decay rate from B-factories:

$$R^* \equiv \frac{BR(B_s^0 \rightarrow D_s^{*-} \mu^+ \nu)}{BR(B^0 \rightarrow D^{*-} \mu^+ \nu)} \sim \frac{|V_{cb}|^2}{BR_{\text{measured B-factories}}}$$


- Result depends on the assumed form factor parametrization:

$$|V_{cb}|_{\text{CLN}} = (41.4 \pm 0.6 (\text{stat}) \pm 0.9 (\text{syst}) \pm 1.2 (\text{ext})) \times 10^{-3}$$

$$|V_{cb}|_{\text{BGL}} = (42.3 \pm 0.8 (\text{stat}) \pm 0.9 (\text{syst}) \pm 1.2 (\text{ext})) \times 10^{-3}$$

- Conclusions:

- First measurement of V_{cb} with pp
- First measurement using B_s^0
- Parametrisation is not responsible for inclusive vs exclusive disagreements
- Result in agreement with the exclusive *and* inclusive averages




New constraints on angle γ

arXiv:2002.08858

- Different yields for B^+ and B^- decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$





New constraints on angle γ

arXiv:2002.08858

■ Different yields for B^+ and B^- decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$

- $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K^0 S K^\pm \pi^\mp$:



$$\begin{aligned}
 N_{SS}^{DK^\pm} &\propto 1 + r_B^2 r_D^2 + 2r_B r_D \kappa_D \cos(\delta_B \pm \gamma - \delta_D) \\
 N_{OS}^{DK^\pm} &\propto r_B^2 + r_D^2 + 2r_B r_D \kappa_D \cos(\delta_B \pm \gamma + \delta_D) \\
 N_{SS}^{D\pi^\pm} &\propto 1 + (r_B^\pi)^2 r_D^2 + 2r_B^\pi r_D \kappa_D \cos(\delta_B^\pi \pm \gamma - \delta_D) \\
 N_{OS}^{D\pi^\pm} &\propto (r_B^\pi)^2 + r_D^2 + 2r_B^\pi r_D \kappa_D \cos(\delta_B^\pi \pm \gamma + \delta_D)
 \end{aligned}$$



	non- K^{*+} region	K^{*+} region
$N_{SS}^{DK^\pm}$	266 ± 27	715 ± 37
$N_{OS}^{DK^\pm}$	336 ± 27	217 ± 22
$N_{SS}^{D\pi^\pm}$	3304 ± 73	8977 ± 106
$N_{OS}^{D\pi^\pm}$	4686 ± 76	3471 ± 66



$$\begin{aligned}
 A_{SS}^{D\pi} &= -0.020 \pm 0.011 \pm 0.003 \\
 A_{OS}^{D\pi} &= 0.007 \pm 0.017 \pm 0.003 \\
 A_{SS}^{DK} &= 0.084 \pm 0.049 \pm 0.008 \\
 A_{OS}^{DK} &= 0.021 \pm 0.094 \pm 0.017
 \end{aligned}$$



New constraints on angle γ

arXiv:1906.08297

■ Different yields for B^0 and \bar{B}^0 decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$

- $B^0 \rightarrow DK^{*0}$ with $D \rightarrow h^\pm h^\mp (h^\pm h^\mp)$:



$$\mathcal{A}_{CP} = \frac{2\kappa r_B^{DK^{*0}} \sin \delta_B^{DK^{*0}} \sin \gamma}{\mathcal{R}_{CP}}$$



Decay channel	\bar{B}^0 yield	B^0 yield
$B^0 \rightarrow D(K^+ K^-) K^{*0}$	67 ± 10	77 ± 11
$B^0 \rightarrow D(\pi^+ \pi^-) K^{*0}$	27 ± 6	40 ± 7
$B^0 \rightarrow D(\pi^+ \pi^- \pi^+ \pi^-) K^{*0}$	32 ± 7	35 ± 8
$B^0 \rightarrow D(K^+ \pi^-) K^{*0}$	786 ± 29	754 ± 29
$B^0 \rightarrow D(\pi^+ K^-) K^{*0}$	76 ± 16	47 ± 15
$B^0 \rightarrow D(K^+ \pi^- \pi^+ \pi^-) K^{*0}$	557 ± 25	548 ± 25
$B^0 \rightarrow D(\pi^+ K^- \pi^+ \pi^-) K^{*0}$	41 ± 14	40 ± 14



$$\begin{aligned} \mathcal{A}_{CP}^{KK} &= -0.05 \pm 0.10 \pm 0.01, \\ \mathcal{A}_{CP}^{\pi\pi} &= -0.18 \pm 0.14 \pm 0.01, \\ \mathcal{A}_{CP}^{4\pi} &= -0.03 \pm 0.15 \pm 0.01, \\ \mathcal{A}_{ADS}^{K\pi} &= 0.047 \pm 0.027 \pm 0.010, \\ \mathcal{A}_{ADS}^{K\pi\pi\pi} &= 0.037 \pm 0.032 \pm 0.010, \end{aligned}$$


CKM angle γ

■ Different yields for B and anti- B decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$
- many $D^{(*)}_{(s)}$ final states:

B decay	D decay	Method	Ref.	Dataset [†]	Status since last combination [3]
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-$	GLW	[14]	Run 1 & 2	Minor update
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-$	ADS	[15]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	GLW/ADS	[15]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-\pi^0$	GLW/ADS	[16]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 h^+h^-$	GGSZ	[17]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 h^+h^-$	GGSZ	[18]	Run 2	New
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 K^+\pi^-$	GLS	[19]	Run 1	As before
$B^+ \rightarrow D^*K^+$	$D \rightarrow h^+h^-$	GLW	[14]	Run 1 & 2	Minor update
$B^+ \rightarrow DK^{*+}$	$D \rightarrow h^+h^-$	GLW/ADS	[20]	Run 1 & 2	Updated results
$B^+ \rightarrow DK^{*+}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	GLW/ADS	[20]	Run 1 & 2	New
$B^+ \rightarrow DK^+\pi^+\pi^-$	$D \rightarrow h^+h^-$	GLW/ADS	[21]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K^+\pi^-$	ADS	[22]	Run 1	As before
$B^0 \rightarrow DK^+\pi^-$	$D \rightarrow h^+h^-$	GLW-Dalitz	[23]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_s^0 \pi^+\pi^-$	GGSZ	[24]	Run 1	As before
$B_s^0 \rightarrow D_s^\mp K^\pm$	$D_s^\pm \rightarrow h^+h^-\pi^\mp$	TD	[25]	Run 1	Updated results
$B^0 \rightarrow D^\mp \pi^\pm$	$D^\pm \rightarrow K^+\pi^-\pi^\pm$	TD	[26]	Run 1	New

[†] Run 1 corresponds to an integrated luminosity of 3 fb⁻¹ taken at centre-of-mass energies of 7 and 8 TeV .
 . Run 2 corresponds to an integrated luminosity of 2 fb⁻¹ taken at a centre-of-mass energy of 13 TeV .





CKM angle γ

- Different yields for B and anti- B decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$
- many $D^{(*)}_{(s)}$ final states:

B decay	D decay	Method	Ref.	Dataset [†]	Status since last combination [3]
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-$	GLW	[14]	Run 1 & 2	Minor update
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-$	ADS	[15]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	GLW/ADS	[15]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow h^+h^-\pi^0$	GLW/ADS	[16]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 h^+h^-$	GGSZ	[17]	Run 1	As before
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 h^+h^-$	GGSZ	[18]	Run 2	New
$B^+ \rightarrow DK^+$	$D \rightarrow K_s^0 K^+\pi^-$	GLS	[19]	Run 1	As before
$B^+ \rightarrow D^*K^+$	$D \rightarrow h^+h^-$	GLW	[14]	Run 1 & 2	Minor update
$B^+ \rightarrow DK^{*+}$	$D \rightarrow h^+h^-$	GLW/ADS	[20]	Run 1 & 2	Updated results
$B^+ \rightarrow DK^{*+}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	GLW/ADS	[20]	Run 1 & 2	New
$B^+ \rightarrow DK^+\pi^+\pi^-$	$D \rightarrow h^+h^-$	GLW/ADS	[21]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K^+\pi^-$	ADS	[22]	Run 1	As before
$B^0 \rightarrow DK^+\pi^-$	$D \rightarrow h^+h^-$	GLW-Dalitz	[23]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_s^0\pi^+\pi^-$	GGSZ	[24]	Run 1	As before
$B_s^0 \rightarrow D_s^\mp K^\pm$	$D_s^\pm \rightarrow h^+h^-\pi^\pm$	TD	[25]	Run 1	Updated results
$B^0 \rightarrow D^\mp\pi^\pm$	$D^\pm \rightarrow K^+\pi^-\pi^\pm$	TD	[26]	Run 1	New


[†] Run 1 corresponds to an integrated luminosity of 3 fb⁻¹ taken at centre-of-mass energies of 7 and 8 TeV .
 . Run 2 corresponds to an integrated luminosity of 2 fb⁻¹ taken at a centre-of-mass energy of 13 TeV .




	γ (°)
LHCb	$74.0^{+5.0}_{-5.8}$
BaBar	69^{+17}_{-16}
World Avg (HFLAV)	$71.1^{+4.6}_{-5.3}$

CKM

1995




2019




- Continuous improvement over the years
- All consistent?

CKM




M. Blanke & A. Buras, EPJ C79 (2019) 159, arXiv:1812.06963
 Emerging ΔM_d Anomaly from Tree-Level Determinations of $|V_{cb}|$ and the Angle γ

- All consistent...?



$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{Bs}}{m_{Bd}} \overset{\text{QCD}}{\xi}^2 \frac{|V_{ts}|^2}{|V_{td}|^2}$$

CKM



M. Blanke & A. Buras, EPJ C79 (2019) 159, arXiv:1812.06963
 Emerging ΔM_d Anomaly from Tree-Level Determinations of $|V_{cb}|$ and the Angle γ

- Interesting $\sim 2\sigma$ tension:

	$\gamma (^{\circ})$
LHCb	$74.0^{+5.0}_{-5.8}$
World Avg (HFLAV)	$71.1^{+4.6}_{-5.3}$
QCD (Δm^{exp} , ξ (Sum Rules))	63.4 ± 0.9

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{Bs}}{m_{Bd}} \overset{\text{QCD}}{\xi}^2 \frac{|V_{ts}|^2}{|V_{td}|^2}$$

On the menu

- Introduction
 - Precision measurements
 - The LHCb physics menu
- Selection of dishes:
 - Recent highlights on CP violation
 - Recent highlights on Rare decays (*aka Flavour Anomalies*)

On the menu

- Introduction

- Precision measurements
- The LHCb physics menu

- Selection of dishes:

- Recent highlights on CP violation
- Recent highlights on Rare decays (*aka Flavour Anomalies*)

- New results

- 1) Lepton flavour non-universality
- 2) Angular analysis of decay
- 3) Search for LFV
- 4) New limit on
- 5) New limit on
- 6) New limit on (x25 !)

$$\Lambda_b^0 \rightarrow p K \mu^+ \mu^-$$

$$B^0 \rightarrow K^{*0} \mu^+ \mu^-$$

$$B^0 \rightarrow K^{*0} \tau^+ \mu^-$$

$$B_s^0 \rightarrow e^+ e^-$$

$$K_S^0 \rightarrow \mu^+ \mu^-$$

$$D_{(s)}^+ \rightarrow h l l'$$




Flavour anomalies

- A remark on consistency

INTERNATIONAL JOURNAL OF HIGH-ENERGY PHYSICS

CERN COURIER


VOLUME 55 NUMBER 9 NOVEMBER 2015



Tensions in the Standard Model


Flavour anomalies? A reminder

- What are the (anomalous) measurements?
 - FCNC: $b \rightarrow sll$
 - LFNU: $b \rightarrow sll$ and $b \rightarrow clv$




FCNC: $b \rightarrow sll$

- $b \rightarrow s$ transition forbidden at tree level in SM



FCNC: $b \rightarrow sll$

- $b \rightarrow s$ transition occurs at loop level
 - Suppressed in SM
 - NP can compete with SM



Flavour-Changing-Neutral-Current-
Electro-Weak-Penguin diagram



FCNC: $b \rightarrow sll$

- $b \rightarrow s$ transition occurs at loop level
 - Suppressed in SM
 - NP can compete with SM

The first penguin:


THE PHENOMENOLOGY OF THE NEXT LEFT-HANDED QUARKS

J. Ellis, M.K. Gaillard ^{*}), D.V. Nanopoulos [†]) and S. Rudaz [‡])
CERN - Geneva



FCNC: $b \rightarrow sll$

- $b \rightarrow s$ transition occurs at loop level
 - LQ quite fashionable these days



deVolkskrant

Moeder aller deeltjes: de zoektocht naar de leptoquark


Is het fundamenteelste deeltje in het universum altijd over het hoofd gezien? Komende week kan de wereld opgeschud worden, als natuurkundigen in Seoul hun resultaten bekendmaken. Leptoquark, onthoud dat woord.

Martijn van Calmthout 29 juni 2018, 11:25

$R_K: B^+ \rightarrow K^+ \mu^+ \mu^-$ and $B^+ \rightarrow K^+ e^+ e^-$

- Similar loop diagram!
- Measure ratio μ/e
- SM expectation: $R_K=1$

$$R_K = \frac{\Gamma(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\Gamma(B^+ \rightarrow K^+ e^+ e^-)}$$




$R_K: B^+ \rightarrow K^+ \mu^+ \mu^- / B^+ \rightarrow K^+ e^+ e^-$

Old result

- Similar loop diagram!
- Measure ratio μ/e
- SM expectation: $R_K=1$


$$R_K = \frac{\Gamma(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\Gamma(B^+ \rightarrow K^+ e^+ e^-)}$$

$$R_K = 0.745^{+0.090}_{-0.074}(\text{stat}) \pm 0.036(\text{syst})$$



LHCb, PRL 113 (2014) 151601

➤ **Lepton flavour
"non-universal" ?**




- Similar loop diagram!
- Measure ratio μ/e
- SM expectation: $R_K=1$


$$R_K = \frac{\Gamma(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\Gamma(B^+ \rightarrow K^+ e^+ e^-)}$$

$$R_K = 0.846^{+0.060}_{-0.054} + 0.016_{-0.014}$$

➤ **Lepton flavour
"non-universal" ?**




LHCb, PRL 122 (2019) 191801



$R_{K^*}: B^0 \rightarrow K^{0*} \mu^+ \mu^-$ and $B^0 \rightarrow K^{0*} e^+ e^-$

- Similar loop diagram!
- Measure ratio μ/e
- SM expectation: $R_{K^*}=1$
- Extra bin at low q^2 ...
 - $q^2 \sim 0$ not helicity suppressed
 - But dominated by photon pole
 - EM coupling to photon undebated...!!



$$R_{K^{*0}} = \begin{cases} 0.66 \pm_{-0.07}^{+0.11} (\text{stat}) \pm 0.03 (\text{syst}) & \text{for } 0.045 < q^2 < 1.1 \text{ GeV}^2/c^4 \\ 0.69 \pm_{-0.07}^{+0.11} (\text{stat}) \pm 0.05 (\text{syst}) & \text{for } 1.1 < q^2 < 6.0 \text{ GeV}^2/c^4 \end{cases}$$

LHCb Coll., JHEP 1708 (2017) 055


- Lepton flavour
"non-universal" ?

$R_{pK}: \Lambda_b^0 \rightarrow pK\mu^+\mu^- / \Lambda_b^0 \rightarrow K^{0*}e^+e^-$


arXiv:1912.08139

- Similar loop diagram!
- Measure ratio μ/e
- SM expectation: $R_{pK}=1$

$$R_{pK}|_{0.1 < q^2 < 6 \text{ GeV}^2/c^4} = 0.86_{-0.11}^{+0.14} \pm 0.05$$



➤ **Lepton flavour
"non-universal" ?**






More LFNU? Semileptonic decays: $b \rightarrow c l \nu$

μ^+ / τ^+

■ $B^0 \rightarrow D^{(*)} l \nu$ Measured ratio τ/μ

- Multiple experiments:
- Multiple c-modes:
- Multiple tau final states:
- Multiple tags:

Belle, BaBar, LHCb
 $D, D^*, J/\psi$
 $\mu, 1\text{-prong}, 3\text{-prong}$
semileptonic, hadronic




and with B_c^+ :


$$\mathcal{R}(J/\psi) = 0.71 \pm 0.17 (\text{stat}) \pm 0.18 (\text{syst})$$

More LFNU? Semileptonic decays: $b \rightarrow c l \nu$

$$\mu^+ / \tau^+$$





- Discrepancy in 2D about 3σ










Decay rates: $b \rightarrow sll$

- Study **same** process with **different** hadrons:



Decay rates: $b \rightarrow sll$


- Decay rate is consistently low:



$$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$$

Old result

- Similar loop diagram!
- More observables
 - Invariant mass of $\mu\mu$ -pair
 - Angles of K and μ




LHCb, JHEP02 (2016) 104, arXiv:1512.04442

$$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$$

Old result

- Similar loop diagram!
- More observables
 - Invariant mass of $\mu\mu$ -pair
 - Angles of K and μ




LHCb, JHEP02 (2016) 104, arXiv:1512.04442

$$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$$

Old result

- Similar loop diagram!
- More observables
 - Invariant mass of $\mu\mu$ -pair
 - Angles of K and μ
- *Many experiments contribute!*

Private compilation, courtesy T. Blake




- LHCb, JHEP02 (2016) 104
- Belle, PRL 118 (2017) 111801
- ATLAS-CONF-2017-023
- CMS, PLB 81 (2018) 517


$$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$$

Old result


- Similar loop diagram!
- More observables
 - Invariant mass of $\mu\mu$ -pair
 - Angles of K and μ
- Debate on SM calculation
 - Non-perturbative “charm loop” effects?



● ATLAS, arXiv:1805.04000
 ● LHCb, JHEP02 (2016) 104
 ◆ Belle, PRL 118 (2017) 111801





or



?

- Updated with (part of) run-2 data




$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$

arXiv:2003.04831


Fit validation

$$\frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d^4(\Gamma + \bar{\Gamma})}{dq^2 d\Omega} \bigg|_P = \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K \right. \\ \left. + \frac{1}{4}(1 - F_L) \sin^2 \theta_K \cos 2\theta_l \right. \\ \left. - F_L \cos^2 \theta_K \cos 2\theta_l + S_3 \sin^2 \theta_K \sin^2 \theta_l \cos 2\phi \right. \\ \left. + S_4 \sin 2\theta_K \sin 2\theta_l \cos \phi + S_5 \sin 2\theta_K \sin \theta_l \cos \phi \right. \\ \left. + \frac{4}{3} A_{FB} \sin^2 \theta_K \cos \theta_l + S_7 \sin 2\theta_K \sin \theta_l \sin \phi \right. \\ \left. + S_8 \sin 2\theta_K \sin 2\theta_l \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_l \sin 2\phi \right]$$

S-wave



Angular acceptance



Systematics

Compatibility


- Run1/2, Magnet polarity, Yields, angular, control channel, ...

Source	F_L	S_3-S_9	P_1-P_8'
Acceptance stat. uncertainty	< 0.01	< 0.01	< 0.01
Acceptance polynomial order	< 0.01	< 0.01	< 0.02
Data-simulation differences	< 0.01	< 0.01	< 0.01
Acceptance variation with q^2	< 0.03	< 0.01	< 0.09
$m(K^+\pi^-)$ model	< 0.01	< 0.01	< 0.01
Background model	< 0.01	< 0.01	< 0.02
Peaking backgrounds	< 0.01	< 0.02	< 0.03
$m(K^+\pi^-\mu^+\mu^-)$ model	< 0.01	< 0.01	< 0.01
$K^+\mu^+\mu^-$ veto	< 0.01	< 0.01	< 0.01
Trigger	< 0.01	< 0.01	< 0.01
Bias correction	< 0.02	< 0.01	< 0.03

$B^0 \rightarrow K^{0*} \mu^+ \mu^-$: more than just P_5'

arXiv:2003.04831


■ Many measurements:



$B^0 \rightarrow K^{0*} \mu^+ \mu^-$: more than just P_5'

arXiv:2003.04831

- Excellent agreement run-1 and 2016:




$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$

arXiv:2003.04831

- What about the tension?


	$4 < q^2 < 6$	$6 < q^2 < 8$	Comb
Run-1	2.8σ	3.0σ	$3.4\sigma^*$
Run-1+2016	2.5σ	2.9σ	3.3σ



- Similar tension in P_5'
- What about overall significance?

Flavour anomalies? Why excitement?

- **Individually**, measurements are consistent with SM
- **Combined** they give an intriguing picture
 - Difference between (lepton) generations?
 - Consistent New Physics scenario possible
 - Simple New Physics scenario possible



On the menu

- Introduction

- Precision measurements
- The LHCb physics menu

- Selection of dishes:

- Recent highlights on CP violation
- Recent highlights on Rare decays (*aka Flavour Anomalies*)

- New results

1) Lepton flavour non-universality

$$\Lambda_b^0 \rightarrow p K \mu^+ \mu^-$$

2) Angular analysis of decay

$$B^0 \rightarrow K^{*0} \mu^+ \mu^-$$

3) Search for LFV

$$B^0 \rightarrow K^{*0} \tau^+ \mu^-$$

4) New limit on

$$B_s^0 \rightarrow e^+ e^-$$


5) New limit on

$$K_S^0 \rightarrow \mu^+ \mu^-$$

6) New limit on (x25 !)



$$D_{(s)}^+ \rightarrow h l l'$$

- A remark on consistency



Intermezzo: Effective couplings

- Historical example





$$\frac{G_F}{\sqrt{2}} = \frac{g^2}{8M_W^2}$$



- Both are correct, depending on the energy scale you consider

Intermezzo: Effective couplings

- Historical example



- Analog: Flavour-changing neutral current





Intermezzo: Effective couplings

- Effective coupling can be of various “kinds”

- Vector coupling
- Axial coupling
- Left-handed coupling (V-A)
- Right-handed (to quarks)
- ...

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{\text{CKM}} \sum_i C_i(\mu) Q_i$$

- Analog: Flavour-changing neutral current



Intermezzo: Effective couplings

■ Effective coupling can be of various “kinds”

- Vector coupling: C_9
- Axial coupling: C_{10}
- Left-handed coupling (V-A): C_9 - C_{10}
- Right-handed (to quarks): C_9', C_{10}', \dots
- Many more! $C_7, C_{1,2}, \dots$

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{\text{CKM}} \sum_i C_i(\mu) Q_i$$



See e.g. Buras & Fleischer, [hep-ph/9704376](https://arxiv.org/abs/hep-ph/9704376)

Semi-Leptonic Operators (fig. 11f):

$$Q_{9V} = (\bar{s}b)_{V-A}(\bar{\mu}\mu)_V$$

$$Q_{10A} = (\bar{s}b)_{V-A}(\bar{\mu}\mu)_A$$

• Analog: Flavour-changing neutral current



Intermezzo: Effective couplings



Model independent fits:

- C_9^{NP} deviates from 0 by $>4\sigma$
- Independent fits by many groups favour:
 - $C_9^{\text{NP}} = -1$ or
 - $C_9^{\text{NP}} = -C_{10}^{\text{NP}}$

➤ All measurements (175) agree with a single (simple?) shift...

LHCb, JHEP 02 (2016) 104

$$\Delta\text{Re}(\mathcal{C}_9) = -1.04 \pm 0.25$$



Intermezzo: Effective couplings



Model independent fits:

- C_9^{NP} deviates from 0 by $>4\sigma$
- Independent fits by many groups favour:
 - $C_9^{\text{NP}} = -1$ or
 - $C_9^{\text{NP}} = -C_{10}^{\text{NP}}$

➤ All measurements (175) agree with a single (simple?) shift...

LHCb, JHEP 02 (2016) 104

$$\Delta\text{Re}(\mathcal{C}_9) = -1.04 \pm 0.25$$



Intermezzo: Effective couplings




Model independent fits:

- C_9^{NP} deviates from 0 by $>4\sigma$
- Independent fits by many groups favour:
 - $C_9^{\text{NP}} = -1$ or
 - $C_9^{\text{NP}} = -C_{10}^{\text{NP}}$

➤ All measurements (175) agree with a single (simple?) shift...


LHCb, JHEP 02 (2016) 104

$$\Delta\text{Re}(C_9) = -1.04 \pm 0.25$$



$B^0 \rightarrow K^{0*} \mu^+ \mu^- : P_5'$

- All (175) measurements favor $C_9^{\text{NP}} = -1.0$
- New P_5' closer to SM, but also in *better* agreement with $C_9^{\text{NP}} = -1.0$
- It is not only about P_5'



Many variables; all sensitive to effective couplings:

- C_7 (photon), C_9 (vector) and C_{10} (axial) couplings hide everywhere:

$$\begin{aligned}
 A_{\perp}^{L,R} &\propto (C_9^{eff} + C_9^{eff'}) \mp (C_{10}^{eff} + C_{10}^{eff'}) \frac{V(q^2)}{m_B + m_{K^*}} + \frac{2m_l}{q^2} (C_7^{eff} + C_7^{eff'}) T_1(q^2) \\
 A_{\parallel}^{L,R} &\propto (C_9^{eff} - C_9^{eff'}) \mp (C_{10}^{eff} - C_{10}^{eff'}) \frac{A_1(q^2)}{m_B + m_{K^*}} + \frac{2m_l}{q^2} (C_7^{eff} - C_7^{eff'}) T_2(q^2) \\
 A_0^{L,R} &\propto (C_9^{eff} - C_9^{eff'}) \mp (C_{10}^{eff} - C_{10}^{eff'}) \times [(m_B^2 - m_{K^*}^2 - q^2)(m_B + m_{K^*} A_1(q^2) - \lambda \frac{A_2(q^2)}{m_B + m_{K^*}})] + \\
 &\quad 2m_l (C_7^{eff} - C_7^{eff'}) [(m_B^2 + 3m_{K^*}^2 - q^2) T_2(q^2) - \frac{\lambda}{m_B^2 - m_{K^*}^2} T_3(q^2)]
 \end{aligned}$$

$$F_L = \frac{A_0^2}{A_{\parallel}^2 + A_{\perp}^2 + A_0^2}$$

$$S_3 = \frac{A_{\perp}^{L2} - A_{\parallel}^{L2}}{A_{\perp}^{L2} + A_{\parallel}^{L2} + A_0^{L2}} + L \rightarrow R$$

$$S_4 = \frac{\Re(A_0^{L*} A_{\parallel}^L)}{|A_0^L|^2 |A_{\parallel}^L|^2 + |A_0^L|^2} + L \rightarrow R$$

$$S_5 = \frac{\Re(A_0^{L*} A_{\perp}^L)}{|A_0^L|^2 + |A_{\perp}^L|^2 + |A_0^L|^2} - L \rightarrow R$$

$$S_6 = \frac{\Re(A_{\perp}^{L*} A_{\parallel}^L)}{|A_{\perp}^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} - L \rightarrow R = \frac{4}{3} A_{FB}$$

$$S_7 = \frac{\Im(A_0^{L*} A_{\parallel}^L)}{|A_0^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} + L \rightarrow R$$




$$S_8 = \frac{\Im(A_0^{L*} A_{\perp}^L)}{|A_0^L|^2 + |A_{\perp}^L|^2 + |A_0^L|^2} + L \rightarrow R$$

$$S_9 = \frac{\Im(A_{\perp}^{L*} A_{\parallel}^L)}{|A_{\perp}^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} - L \rightarrow R$$

$$\begin{aligned}
 \frac{1}{\Gamma} \frac{d^3(\Gamma + \bar{\Gamma})}{d \cos \theta_{\ell} d \cos \theta_K d \phi} &= \frac{9}{32\pi} \left[\frac{3}{4} (1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K + \frac{1}{4} (1 - F_L) \sin^2 \theta_K \cos 2\theta_{\ell} \right. \\
 &\quad \left. - F_L \cos^2 \theta_K \cos 2\theta_{\ell} + \right. \\
 &\quad S_3 \sin^2 \theta_K \sin^2 \theta_{\ell} \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_{\ell} \cos \phi + \\
 &\quad S_5 \sin 2\theta_K \sin \theta_{\ell} \cos \phi + S_6 \sin^2 \theta_K \cos \theta_{\ell} + \\
 &\quad S_7 \sin 2\theta_K \sin \theta_{\ell} \sin \phi + \\
 &\quad \left. S_8 \sin 2\theta_K \sin 2\theta_{\ell} \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_{\ell} \sin 2\phi \right]
 \end{aligned}$$


Best fit

- Improved fit for $C_9^{\text{NP}} = -1.0$



Emerging patterns of New Physics with and without Lepton Flavour Universal contributions

Marcel Algueró^{a,b}, Bernat Capdevila^{a,b,c}, Andreas Crivellin^{d,e}, Sébastien Descotes-Genon^f, Pere Masjuan^{a,b}, Joaquim Matias^{a,b}, Martín Novoa Brunet^f and Javier Virto^g.



- There is a reduction of the internal tensions between some of the most relevant observables of the fit, in particular, between the new averages of R_K and P'_5 . This leads to an increase in consistency between the different anomalies. This is illustrated

1D Hyp.	All			
	Best fit	$1\sigma/2\sigma$	Pull _{SM}	p-value
$C_{9\mu}^{NP}$	-1.03	$[-1.19, -0.88]$ $[-1.33, -0.72]$	6.3	37.5 %
$C_{9\mu}^{NP} = -C_{10\mu}^{NP}$	-0.50	$[-0.59, -0.41]$ $[-0.69, -0.32]$	5.8	25.3 %

- The reduced uncertainties of the $B \rightarrow K^* \mu \mu$ data and its improved internal consistency sharpen statistical statements on the hypotheses considered. There is a significant increase of the statistical exclusion of the SM hypothesis as its p-value is reduced down to 1.4% (i.e. 2.5σ). The Pull_{SM} of the 6D fit is now higher (5.8σ).

arXiv:1903.09578, **addendum 6 Apr 2020**


- Similar picture as before
- Reduction of internal tensions
- Increase of statistical exclusion of SM hypothesis
 - p-value 1.4%, Pull 5.8σ

Outlook

2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034
		Run III						Run IV					Run V	
LS2	?					LS3					LS4			
LHCb 40 MHz UPGRADE I		$L = 2 \times 10^{33}$			LHCb Consolidate: Upgr Ib			$L = 2 \times 10^{33}$ 50 fb^{-1}			LHCb UPGRADE II		$L=1-2 \times 10^3$ 300 fb^{-1}	
ATLAS Phase I Upgr		$L = 2 \times 10^{34}$			ATLAS Phase II UPGRADE			HL-LHC $L = 5 \times 10^{34}$			ATLAS		HL-LHC $L = 5 \times 10^3$	
CMS Phase I Upgr		300 fb^{-1}			CMS Phase II UPGRADE						CMS		3000 fb^{-1}	
Belle II		5 ab^{-1}	$L = 8 \times 10^{35}$	50 ab^{-1}				LHC schedule: Frederick Bordry, 2019 https://lhc-commissioning.web.cern.ch/schedule/LHC-long-term.htm						



Conclusions

- Precision measurements to scrutinize the Standard Model
- Precision measurements reach very high mass scales
- Precision measurements are not yet precise enough




2019



C_{10}^{NP} vs C_9^{NP}



2030



CKM: γ , V_{ub} , Δm_s



What NP could it be?

- If interpreted as NP signals, both set of anomalies are not in contradiction among themselves & with existing low- & high-energy data.
Taken together, they point out to NP coupled mainly to 3rd generation, with a flavor structure connected to that appearing in the SM Yukawa couplings


G. Isidori, *Implications workshop*, CERN, 10 Nov 2017
<https://indico.cern.ch/event/646856/timetable/>

- Anomalous measurements:
 - FCNC: $b \rightarrow sll$
 - LFNU: $b \rightarrow sll$ and $b \rightarrow clv$
- What are the interpretations?


Model building

- Most popular models: Z' or Leptoquark





SM



$SU(2)'$




Leptoquark




Model building

Step 1: Effective theory




$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} - \frac{1}{v^2} \lambda_{ij}^q \lambda_{\alpha\beta}^\ell \left[C_T (\bar{Q}_L^i \gamma_\mu \sigma^a Q_L^j) (\bar{L}_L^\alpha \gamma^\mu \sigma^a L_L^\beta) + C_S (\bar{Q}_L^i \gamma_\mu Q_L^j) (\bar{L}_L^\alpha \gamma^\mu L_L^\beta) \right]$$

Observable	Experimental bound	Linearised expression
$R_{D^{(*)}}^{\tau\ell}$	1.237 ± 0.053	$1 + 2C_T(1 - \lambda_{sb}^q V_{tb}^*/V_{ts}^*)(1 - \lambda_{\mu\mu}^\ell/2)$
$\Delta C_9^\mu = -\Delta C_{10}^\mu$	-0.61 ± 0.12 [36]	$-\frac{\pi}{\alpha_{\text{em}} V_{tb} V_{ts}^*} \lambda_{\mu\mu}^\ell \lambda_{sb}^q (C_T + C_S)$
$R_{b \rightarrow c}^{\mu e} - 1$	0.00 ± 0.02	$2C_T(1 - \lambda_{sb}^q V_{tb}^*/V_{ts}^*) \lambda_{\mu\mu}^\ell$
$B_{K^{(*)}\nu\bar{\nu}}$	0.0 ± 2.6	$1 + \frac{2}{3} \frac{\pi}{\alpha_{\text{em}} V_{tb} V_{ts}^* C_{\nu}^{\text{SM}}} (C_T - C_S) \lambda_{sb}^q (1 + \lambda_{\mu\mu}^\ell)$
$\delta g_{\tau_L}^Z$	-0.0002 ± 0.0006	$0.033C_T - 0.043C_S$
$\delta g_{\nu_\tau}^Z$	-0.0040 ± 0.0021	$-0.033C_T - 0.043C_S$
$ g_\tau^W/g_\ell^W $	1.00097 ± 0.00098	$1 - 0.084C_T$
$\mathcal{B}(\tau \rightarrow 3\mu)$	$(0.0 \pm 0.6) \times 10^{-8}$	$2.5 \times 10^{-4} (C_S - C_T)^2 (\lambda_{\tau\mu}^\ell)^2$



Step 2: Simplified models




$SU(2)_L$ -singlet vector leptoquark, $U_1^\mu \equiv (\mathbf{3}, \mathbf{1}, 2/3)$

$$\begin{aligned} \mathcal{L}_U &= -\frac{1}{2} U_{1,\mu\nu}^\dagger U^{1,\mu\nu} + M_U^2 U_{1,\mu}^\dagger U_1^\mu + g_U (J_U^\mu U_{1,\mu} + \text{h.c.}) \\ J_U^\mu &\equiv \beta_{i\alpha} \bar{Q}_i \gamma^\mu L_\alpha \end{aligned}$$


Model building

- Many models! See e.g.:

➤ Possible BSM models




- Heavy Z' model
- $SU(2)_L$ singlet or triplet
- arXiv:1403.1269, 1501.00993, 1503.03477, 1611.02703, ...




- Leptoquark model
- Spin 0 or 1
- arXiv:01511.01900, 1503.01084, 1704.05835, 1512.01560, 1511.06024, 1408.1627, ...

arXiv:1706.07808




- Other new heavy scalars/vectors also leptoquark possible
- arXiv:01509.05020, 1608.07832, 1704.05438, 1607.01659, 1704.07845, hep-ph/0610037, ...



Model building

Ingredients



- NP: large coupling $b \rightarrow c \tau \nu$
 - Large coupling to 3rd gen leptons
 - Left-handed coupling (no RH neutrino)
- NP: small (non-vanishing) coupling $b \rightarrow s \mu \mu$
 - Small coupling to 2nd gen leptons
 - Left-handed coupling (from C_9)



Model building

Ingredients

- NP: large coupling $b \rightarrow c \tau \nu$
 - Large coupling to 3rd gen leptons
 - Left-handed coupling (no RH neutrino)
- NP: small (non-vanishing) coupling $b \rightarrow s \mu \mu$
 - Small coupling to 2nd gen leptons
 - Left-handed coupling (from C_9)



G.Isidori

J.M.Camalich

Experimental constraints

- High p_T searches (No $\pi\pi$ resonance: no s-channel Z')
- Radiative constr. $\tau \rightarrow \mu \nu \nu$
- B_s^0 mixing (No tree level NP: small b_s implies large $\tau \nu$)
- B_c^+ lifetime (Scalar LQ increases $\text{BR}(B_c^+ \rightarrow \tau^+ \nu)$)

Vector LQ favoured
over
Scalar LQ or Z'

Model building

$SU(2)_L$ -singlet vector leptoquark

emerges as a particularly simple and successful framework.

- Many more experimental handles; predictions can be checked!

- Universal for all $b \rightarrow c \tau \nu$:
 - Accurate $R(D^*)$, $R(J/\psi)$, ...

$$\frac{R_D}{(R_D)_{SM}} = \frac{\Gamma(B \rightarrow D^* \tau \nu) / \Gamma_{SM}}{\Gamma(B \rightarrow D^* \mu \nu) / \Gamma_{SM}} = \frac{\Gamma(B_c \rightarrow \psi \tau \nu) / \Gamma_{SM}}{\Gamma(B_c \rightarrow \psi \mu \nu) / \Gamma_{SM}} = \frac{\Gamma(\Lambda_b \rightarrow \Lambda_c \tau \nu) / \Gamma_{SM}}{\Gamma(\Lambda_b \rightarrow \Lambda_c \mu \nu) / \Gamma_{SM}} = \dots$$

- Strong coupling to τ au's:
 - Measure e.g. $B^0 \rightarrow K^* \tau \tau$

- LFNU linked with LFV:
 - Look for e.g. $B^0 \rightarrow K^* \tau \mu$
 - $BR(\tau \rightarrow \mu \mu \mu) \sim 10^{-9}$

	$\mu\mu$ (ee)	$\tau\tau$	$\nu\nu$	$\tau\mu$	μe
$b \rightarrow s$	R_K, R_{K^*} O(20%)	$B \rightarrow K^{(*)} \tau\tau$ $\rightarrow 100 \times SM$	$B \rightarrow K^{(*)} \nu\nu$ O(1)	$B \rightarrow K \tau\mu$ $\rightarrow \sim 10^{-6}$	$B \rightarrow K \mu e$???
$b \rightarrow d$	$B_d \rightarrow \mu\mu$ $B \rightarrow \pi \mu\mu$ $B_s \rightarrow K^{(*)} \mu\mu$ O(20%) [$R_K = R_\pi$]	$B \rightarrow \pi \tau\tau$ $\rightarrow 100 \times SM$	$B \rightarrow \pi \nu\nu$ O(1)	$B \rightarrow \pi \tau\mu$ $\rightarrow \sim 10^{-7}$	$B \rightarrow \pi \mu e$???

- c, u symmetry:
 - Study suppressed semileptonic

$$\frac{\Gamma(B \rightarrow \pi \tau \nu) / \Gamma_{SM}}{\Gamma(B \rightarrow \pi \mu \nu) / \Gamma_{SM}} = \frac{\Gamma(\Lambda_b \rightarrow p \tau \nu) / \Gamma_{SM}}{\Gamma(\Lambda_b \rightarrow p \mu \nu) / \Gamma_{SM}} = \frac{\Gamma(B_s \rightarrow K^* \tau \nu) / \Gamma_{SM}}{\Gamma(B_s \rightarrow K^* \mu \nu) / \Gamma_{SM}} = \dots = \frac{R_D}{(R_D)_{SM}}$$


- B_s mixing
 - O(1-10%) effect on Δm_s

Buttazzo, Greife, Isidori, Marzocca, B-physics anomalies: a guide to combined explanations, JHEP 1711 (2017) 044

Model building


- Many more experimental handles; predictions can be checked!
- High p_T signatures?

– LQ pairs





– LQ t-channel in $bb \rightarrow \tau\tau$

Reachable
during HL-LHC



– Single production channel
(dominant?)



The need for more precision

Imagine if Fitch and Cronin had stopped at the 1% level, how much physics would have been missed”

– A.Soni

- “A special search at Dubna was carried out by Okonov and his group. They did not find a single $K_L^0 \rightarrow \pi^+ \pi^-$ event among 600 decays into charged particles (Anikira et al., JETP 1962). At that stage the search was terminated by the administration of the lab. The group was unlucky.”

– L.Okun

(remember: $B(K_L^0 \rightarrow \pi^+ \pi^-) \sim 2 \cdot 10^{-3}$)